




Distinguishing science from pseudoscience in commercial respiratory interventions: an evidence-based guide for health and exercise professionals

Camilla R. Illidi¹ · Lee M. Romer² · Michael A. Johnson³ · Neil C. Williams³ · Harry B. Rossiter⁴ · Richard Casaburi⁴ · Nicholas B. Tiller⁴ 

Received: 23 December 2022 / Accepted: 19 February 2023

© The Author(s), under exclusive licence to Springer-Verlag GmbH Germany, part of Springer Nature 2023, corrected publication 2023

Abstract

Respiratory function has become a global health priority. Not only is chronic respiratory disease a leading cause of worldwide morbidity and mortality, but the COVID-19 pandemic has heightened attention on respiratory health and the means of enhancing it. Subsequently, and inevitably, the respiratory system has become a target of the multi-trillion-dollar health and wellness industry. Numerous commercial, respiratory-related interventions are now coupled to therapeutic and/or ergogenic claims that vary in their plausibility: from the reasonable to the absurd. Moreover, legitimate and illegitimate claims are often conflated in a wellness space that lacks regulation. The abundance of interventions, the range of potential therapeutic targets in the respiratory system, and the wealth of research that varies in quality, all confound the ability for health and exercise professionals to make informed risk-to-benefit assessments with their patients and clients. This review focuses on numerous commercial interventions that purport to improve respiratory health, including nasal dilators, nasal breathing, and systematized breathing interventions (such as pursed-lips breathing), respiratory muscle training, canned oxygen, nutritional supplements, and inhaled L-menthol. For each intervention we describe the premise, examine the plausibility, and systematically contrast commercial claims against the published literature. The overarching aim is to assist health and exercise professionals to distinguish science from pseudoscience and make pragmatic and safe risk-to-benefit decisions.

Keywords Asthma · COPD · Exercise · Disease · Lung function · Nutrition · Pulmonary

Communicated by Michael I Lindinger.

✉ Nicholas B. Tiller
nicholas.tiller@lundquist.org

¹ Clinical Exercise and Respiratory Physiology Laboratory, Department of Kinesiology and Physical Education, Faculty of Education, McGill University, Montréal, QC, Canada

² Division of Sport, Health and Exercise Sciences, College of Health, Medicine and Life Sciences, Brunel University London, Uxbridge, UK

³ Exercise and Health Research Group, Sport, Health and Performance Enhancement (SHAPE) Research Centre, School of Science and Technology, Nottingham Trent University, Nottingham, Nottinghamshire, UK

⁴ Institute of Respiratory Medicine and Exercise Physiology, Division of Respiratory and Critical Care Physiology and Medicine, The Lundquist Institute for Biomedical Innovation at Harbor-UCLA Medical Center, 1124 W. Carson Street, CDCRC Building, Torrance, CA 90502, USA

Introduction

The human respiratory system comprises the upper respiratory tract (nasal and oral cavities, pharynx, and larynx), lower respiratory tract (trachea, and bronchial tree), lung parenchyma, pulmonary vasculature, and respiratory muscles (e.g., diaphragm, abdominals, intercostals). This coordinated arrangement, under neural control from central and peripheral chemoreceptors and respiratory centers in the brain, transfers oxygen from the atmosphere to the pulmonary circulation and carbon dioxide in the opposite direction. With a few notable exceptions (e.g., high-intensity exercise, hypoxic environments), the healthy respiratory system is unlikely to present a significant limitation to gas exchange or O₂ transport (Dempsey et al. 2020).

In recent years, the respiratory system has become a target of the multi-trillion-dollar commercial health and wellness industry. Therein, numerous respiratory-related products and strategies (e.g., respiratory muscle training

devices, nasal strips, deep breathing regimens) are sold to the consumer alongside therapeutic and/or ergogenic claims that vary in their plausibility: from the reasonable (mitigate stress, improve perceptions, improve lung and respiratory muscle function); to the questionable (increase oxygen transport, “boost” immune function); to the absurd (increase “energy flow” and promote healing). Furthermore, due to lax regulations in the wellness space and little obligation for marketing to conform to scientific or ethical standards, it is common for legitimate and illegitimate claims to be conflated (Tiller et al. 2022). The current ‘wellness’ paradigm thus makes it difficult for health and exercise professionals to make informed risk-to-benefit assessments with their patients and clients.

Several factors underpin the accelerating commercial popularity of respiratory-related interventions, the most pertinent being the COVID-19 pandemic which has heightened attention on respiratory health and potential means of enhancing it. But even before COVID-19, chronic respiratory disease (such as chronic obstructive pulmonary disease [COPD]) was a leading cause of morbidity and mortality (World Health Organization 2022), conferring a considerable and growing economic burden (Ehteshami-Afshar et al. 2016). Respiratory disease has also received growing coverage in the media owing to the pressing issue of climate change and worsening air quality (Barnes et al. 2013). Respiratory function has thus become a global health priority. To compound the problem, respiratory physiology is a complex discipline that is poorly understood by the public, and its mechanisms can thus be easily misappropriated for commercial gain.

This review examines commercial interventions that purport to influence aspects of the respiratory system to improve respiratory function, respiratory health, and/or exercise responses. The interventions selected for inclusion were nasal dilators, nasal breathing, and systematized breathing interventions, respiratory muscle training, canned oxygen, nutritional supplements, and inhaled L-menthol. For each intervention we describe its premise, examine its plausibility, and contrast commercial claims against the published literature. The overarching aim of this paper is to provide an evidence-based guide for health and exercise professionals—to help them distinguish science from pseudoscience in respiratory physiology and assist them in making safe and pragmatic risk-to-benefit decisions.

Methods

In January 2022, the first and corresponding authors (CRI and NBT, respectively) convened a meeting of recognized experts in the fields of respiratory medicine and exercise physiology. After several rounds of discussion, all authors agreed that the products/strategies to be included should

be non-medical, commercial interventions, and excluded if they were controlled drugs and/or regulated by the FDA as “medical devices”. The list was not exhaustive but limited to the interventions most prevalent in the health and wellness industry and that were coupled to the most conspicuous claims. A list of commercial claims was then compiled from websites, press releases, and relevant media, after which peer-reviewed articles were searched via PubMed (no date restriction). The literature search-terms comprised the relevant intervention (e.g., nasal dilators, respiratory muscle training, etc.) alongside various combinations of the following: breathlessness; dyspnea; lung; lung function; pulmonary; respiratory; respiratory function; respiratory health; respiratory symptoms; pathophysiology. All article types—meta-analyses, systematic reviews, randomized-controlled trials (RCTs), exploratory studies, confirmatory studies, and case reports—were included, and the reference lists of articles selected for inclusion were manually searched for additional literature. The manuscript was drafted, and after several rounds of discussion and refinement, all authors agreed upon the evidence summaries and recommendations and approved the final work.

Evidence review

Nasal dilators

Premise and plausibility

External nasal dilators (ENDs) are applied horizontally to the skin of the nasal dorsum whereas internal nasal dilators (INDs) are placed inside the nostrils. Both purportedly increase nostril patency by preventing the nasal wings from collapsing during inspiration (Dinardi et al. 2014). The devices were originally developed to aid with sleep-related issues (e.g., snoring and apnea) but their widespread use at the Atlanta Olympic Games in 1996 made them popular with exercisers and athletes (Dinardi et al. 2014). Using magnetic resonance imaging, Bishop et al. (2016) showed that an END (Breathe Right[®]) significantly enlarged the anterior nasal passage compared to a placebo. Using acoustic rhinometry, Griffin et al. (1997) observed increased nasal valve area with the same device. Although ENDs had no effect on plethysmography-derived measures of nasal resistance (Vermoen et al. 1998) or maximum expiratory flows (Di Somma et al. 1999) in healthy individuals, others have shown that ENDs increased nasal inspiratory flow during normal and forced breathing (Vermoen et al. 1998; Di Somma et al. 1999). Thus, ENDs likely improve nostril patency by supporting the lateral nasal vestibular walls, manifesting as a slight increase in inspiratory nasal flow at rest and during maximal inspiratory maneuvers. The bulk of literature has

focused on whether there is any subsequent clinical or ergogenic benefit.

Literature

Articles were excluded if nasal dilators were simultaneously applied with other breathing interventions. Most studies on nasal dilators evaluated their effect on sleep-related issues including sleep quality, snoring, and obstructive sleep apnea, generally showing subjective (but not objective) outcomes. For example, using an END, Wenzel et al. (1997) reported improved subjective ratings of nose breathing at rest but no changes in objective (polysomnography-derived) measures of obstructive sleep apnea. Similarly, several non-placebo-controlled studies showed improved subjective ratings of sleep quality (e.g., insomnia severity, sleep-disordered breathing) and quality of life with ENDs (Krakow et al. 2006; Gelardi et al. 2019). When an IND (Nas-Air[®]) and an END (Breathe Right[®]) were compared, the former conferred better subjective ratings of sleep quality (assessed via visual analogue scale) (Gelardi et al. 2019). One placebo-controlled study in patients with upper-airway resistance syndrome found that desaturation time during sleep (the percentage of time that SpO₂ was > 2% below waking values) was significantly lower with an END (Breathe Right[®]) versus placebo (9.1 vs. 12.2%). However, there were no other effects on cardiorespiratory variables, sleep architecture, or sleep latency (Bahammam et al. 1999).

Data on snoring are equivocal. Gelardi et al. (2019) found that snoring time was reduced with both an IND and END, whereas Wenzel et al. (1997) showed that ENDs had no effect on the frequency of snoring events. When healthy subjects with nasal congestion were randomized to an END group (Breathe Right[®]) or a placebo group for two weeks, both devices equally improved subjective ratings of sleep quality and subjective ratings of nasal congestion, suggesting a potent placebo effect (Noss et al. 2019). Lastly, Sadan et al. (2005) showed that nasal dilators, when used by females during childbirth, improved subjective ratings of “ease of breathing” but had no effect on objective markers of labor progression or recovery. Collectively, the data suggest that nasal dilators (mainly ENDs) may improve perceptions of nasal breathing and subjective ratings of sleep quality. However, they are unlikely to influence objective markers of obstructive sleep apnea (Camacho et al. 2016) and the data on snoring frequency and duration are equivocal. In fact, several authors have proposed ENDs as an effective placebo intervention in RCTs that explore treatment options in obstructive sleep apnea (Amaro et al. 2012; Yagihara et al. 2017).

On the premise that ENDs increase some measures of nasal patency and nasal inspiratory flow, their potential to

improve exercise capacity has also been explored. Despite a few reports of favorable outcomes (Griffin et al. 1997; Dinardi et al. 2013, 2017), a recent systematic review and meta-analysis of 19 articles concluded that ENDs elicited “no improvement in $\dot{V}O_2$ max, HR and RPE outcomes in healthy individuals during [maximal or submaximal] exercise” (Dinardi et al. 2021). Other studies, using esophageal balloon catheters, showed no effect of ENDs on inspiratory elastic work, inspiratory resistive work, or expiratory resistive work during submaximal or maximal exercise (O’Kroy et al. 2001). There was also no effect of ENDs on recovery of $\dot{V}O_2$, \dot{V}_E , or HR after exercise when compared to a placebo or a no-intervention control (Thomas et al. 2001), and no effect of ENDs on blood lactate responses after exercise in sedentary or endurance trained women (Boggs et al. 2008). Thus, nasal dilators appear to have no meaningful influence on physiological variables during or after exercise.

The commercial claims of one specific IND (Turbine[™]) have been scrutinized by several studies that showed no benefit on respiratory mechanics or exercise tolerance. One such study, a sham-controlled trial using esophageal balloon catheters to measure respiratory mechanics during incremental cycle ergometry, showed that the IND did not reduce the work of breathing and had no effect on exertional dyspnea or exercise capacity (Schaeffer et al. 2021). Another RCT that evaluated the effect of the Turbine[™] on 20-km cycling time-trial performance showed no influence on mean power output (Adams and Peiffer 2017). Favorable data from a clinical trial posted on the manufacturer’s website have not been peer reviewed or published at the time of this writing.

Evidence summary and recommendations

Primary outcomes from the literature on nasal dilators are summarized in Fig. 1. Both ENDs and INDs enlarge and stabilize the nasal valves thereby mitigating their collapse during high flow inspiration and increasing inspiratory nasal airflow during maximal inspiratory efforts. Current evidence suggests that nasal dilators may improve perceptions of nasal breathing and subjective ratings of sleep quality but are unlikely to influence objective markers of obstructive sleep apnea. The data on snoring frequency and duration are equivocal. Most studies show no effect of nasal dilators on cardiorespiratory function or ratings of perceived exertion during exercise, no effect on exercise capacity, and no effect on physiological variables during the acute phase of recovery.




Fig. 1 Primary outcomes from the literature on internal and external nasal dilators. ↑ evidence of increase, ↓ evidence of decrease, ↔ evidence of no change, *IND* internal nasal dilator, *END* external nasal dilator,

OSA obstructive sleep apnea, $\dot{V}O_2$ max maximal oxygen uptake, *HR* heart rate, *RPE* ratings of perceived exertion

Nasal breathing

Premise and plausibility

In humans, nitric oxide (NO) is a vasodilator (Morris and Rich 1997) and mild bronchodilator (Kacmarek et al. 1996) that was first identified in expired gas in the 1990s (Gustafsson et al. 1991). Functionally, the two NO isoforms are “constitutive” and “inducible” NO, with most being produced in the paranasal sinuses (Ricciardolo 2003). In fact, the paranasal sinuses produce considerably greater amounts of NO than either the mouth or the trachea [56 vs. 14 vs. 6 ppb, respectively; (Törnberg et al. 2002)]. It has been suggested that nasally-derived NO can evoke airway smooth muscle relaxation, inhibit smooth muscle proliferation, and protect against excessive bronchoconstriction (Ricciardolo 2003). Others suggest that nasal breathing might attenuate pulmonary hypertension by vasodilating the pulmonary vasculature (Settergren et al. 1998). Although exogenous (supplementary) NO is known to reduce vascular resistance and increase pulmonary blood flow in healthy and patient populations (Settergren et al. 1998; Crespo et al. 2010), the concentration

of endogenous (nasally-derived) NO is considerably lower than the concentrations used in NO-enriched air (Törnberg et al. 2002). Therefore, an important consideration is whether increased NO uptake via nasal breathing exerts meaningful effects in healthy or patient populations.

Literature

Articles were excluded if they reported on exogenous (supplementary) NO inhalation, if they studied exhaled NO as a tool for assessing airway inflammation, or if nasal breathing was studied in combination with other breathing interventions (e.g., deep/slow breathing). Using single photon emission computed tomography during separate bouts of upright nasal or oral breathing in healthy adults, Crespo et al. (2010) found that nasal breathing elicited blood flow redistribution from caudal and dorsal regions of the lung to the less-perfused cranial and ventral regions. For the poorly perfused lung regions, such as the apical region, this represented a 24% increase in blood flow. Similar effects were observed when exogenous NO mixtures were inhaled orally, supporting the

hypothesis that blood flow redistribution had been mediated by NO. Others showed that oxygen tension across the chest wall (assessed using transcutaneous electrodes) was increased in healthy subjects during nasal breathing versus oral breathing (Lundberg et al. 1996); however, the effects were very small and the clinical significance thus unclear.

Limited data also suggest a possible therapeutic benefit of nasal breathing in patients with respiratory disease. For example, intubated patients who were unable to rebreathe their own nasally-derived NO exhibited an 18% increase in PaO₂, and an 11% decrease in pulmonary vascular resistance index, when gas derived from the patient's nose was aspirated and fed into the inspiratory limb of the ventilator (Lundberg et al. 1996). Although the exact mechanism was unclear, the authors postulated that sinus-derived NO may act as an “aerocrine messenger” that selectively dilates vessels supplying well-ventilated areas of the lung. Pulmonary vascular resistance also decreased in patients recovering from thoracic surgery when they engaged in nasal versus oral breathing, although there was no difference in O₂ and CO₂ partial pressures of arterial and mixed venous blood between the two breathing techniques (Settergren et al. 1998). Pertinently, exogenous NO inhaled orally at “nasal physiologic concentrations” of 10–100 ppb evoked pulmonary vasodilatation and improved pulmonary gas exchange in patients with acute respiratory distress syndrome (Mourgeon et al. 1997) and acute respiratory failure (Gerlach et al. 1993). Collectively, these data support the notion that increasing NO uptake via nasal breathing may provide clinically meaningful benefits in certain patient populations.

The potential benefit of nasal breathing at rest has led to the suggestion that it may improve physiological responses to exercise. However, the feasibility of nasal breathing during exercise should first be considered. Healthy adults spontaneously switch from nasal to oronasal breathing at minute ventilations of 35–45 L·min⁻¹ (Niinimaa et al. 1980; Becquemin et al. 1991; Bennett et al. 2003). Moreover, without prior habituation, healthy adults can only maintain nasal breathing up to ~80% $\dot{V}O_2$ max (LaComb et al. 2017). Nevertheless, when preceded by an extensive training period (> 6 months), nasal breathing may be feasible during high-intensity and even maximal exercise without compromising $\dot{V}O_2$ max (Hostetter et al. 2016; Dallam et al. 2018). This leads to the separate question of efficacy: does nasal breathing during exercise provide any physiological advantage over oral or oronasal breathing?

In a mixed-sex cohort of healthy adults, LaComb et al. (2017) showed that nasal breathing elicited lower $\dot{V}O_2$, $\dot{V}CO_2$, and \dot{V}_E at given submaximal exercise intensities (50%, 65%, and 80% of treadmill-derived $\dot{V}O_2$ max) when compared to oral breathing, although the physiological mechanism was unclear. A possible limitation of the study was that exercise bouts lasted only 4 min; steady state responses may take considerably longer,

particularly in an untrained cohort with a slow kinetic response. The authors also concluded that, when all variables were considered together, “it is likely that oral breathing represents the more efficient mode [of breathing], particularly at higher exercise intensities”. In another study, 10 healthy subjects who were habituated to nasal breathing exhibited lower ventilatory equivalents for O₂ and CO₂ during nasal-only exercise versus oral-only exercise (differences mediated primarily by significantly lower \dot{V}_E), without a change in $\dot{V}O_2$ max or time to exhaustion (Dallam et al. 2018). It is unclear from these studies whether the physiological responses to nasal breathing were derived from increased NO uptake or another mechanism. It is possible that healthy subjects have a blunted ventilatory response at maximal exercise with nasal breathing owing to attenuated tidal volumes and respiratory frequencies (Morton et al. 1995). This may partly explain greater end-tidal CO₂ partial pressure (P_{ET}CO₂) during nasal versus oral breathing, both at rest and during submaximal exercise (Tanaka et al. 1988; Dallam et al. 2018).

Although oral and nasal breathing evoke similar ratings of perceived exertion during exercise, nasal breathing may result in an “unacceptable sensation of air hunger” (Hostetter et al. 2016; Dallam et al. 2018) possibly owing to the development of hypercapnia (Banzett et al. 2021). In turn, increased air hunger has the potential to alter breathing patterns and reduce exercise tolerance (Dallam et al. 2018). In patients with COPD who report pre-existing sensations of “unsatisfied inspiration” (Philips et al. 2021), additional air hunger evoked by nasal breathing could exacerbate respiratory symptoms. Two early studies in patients with asthma ($n=5$ and $n=12$, respectively) found that nasal breathing during exercise reduced the incidence and/or severity of post-exercise bronchoconstriction relative to oral breathing (Shturman-Ellstein et al. 1978; Mangla and Menon 1981). However, it is unclear whether these findings were exclusively the result of greater NO uptake or were also influenced by an increased humidity of inspired air (Naclerio et al. 2007) which has also been shown to mitigate the severity of exercise-induced bronchoconstriction (Anderson and Kippelen 2012). In any case, these data support the hypothesis that the nasopharynx and oropharynx play an important role in mediating exercise-induced bronchoconstriction.

Evidence summary and recommendations

Primary outcomes from the literature on nasal breathing are summarized in Fig. 2. Data suggest that nasal breathing may improve arterial oxygenation in critically ill patients at rest, but there is little evidence that such benefits extend to healthy subjects. Nasal breathing is feasible during submaximal exercise and even maximal exercise after extensive habituation, but there is little-to-no data supporting a subsequent benefit on exercise capacity in healthy individuals. There is some evidence of reduced prevalence and/or




Fig. 2 Primary outcomes from the literature on nasal breathing. ↑ evidence of increase; ↓ evidence of decrease; ↔ evidence of no change, NO nitric oxide, $\dot{V}O_2$ oxygen uptake, \dot{V}_E minute ventilation, $\dot{V}_E/\dot{V}O_2$ ventilatory equivalent for oxygen, $\dot{V}_E/\dot{V}CO_2$ ventilatory equivalent for carbon

dioxide, $P_{ET}CO_2$ end-tidal partial pressure of carbon dioxide, f_R respiratory frequency, V_T tidal volume, $\dot{V}O_2$ max maximal oxygen uptake

severity of post-exercise bronchoconstriction with nasal breathing, but due to potential hypoventilation and increased sensations of “air hunger”, nasal breathing during exercise is not recommended for COPD patients. Its use in patients with other respiratory diseases should be considered on a case-by-case basis.

Systematized breathing strategies

Premise and plausibility

Breathing interventions generally comprise one-or-more of the following techniques: nasal inspiration, preferential activation of the diaphragm during inspiration, deep/slow breaths, breath-hold at end-inspiratory lung volume, prolonged expiration, and expiration through pursed lips. Most breathing interventions encourage inspiration through the nose. This approach may increase the uptake of NO (see “Nasal breathing”) and warm/humidify the inspired air (Naclerio et al. 2007). Inspiration and expiration are usually required to be deep and slow to increase tidal volume and extend the respiratory cycle (Ubolnuar et al. 2019). A prolonged expiration may also help decrease expiratory reserve volume which, in turn, may partially mitigate air

trapping and dynamic hyperinflation, and reduce respiratory symptoms in certain populations (e.g., COPD). Deep/slow breathing, particularly interventions with prolonged expiration, have also been shown to increase heart rate variability and respiratory sinus arrhythmia through mediating effects on the parasympathetic nervous system (Zaccaro et al. 2018). This is an expanding area of research. Indeed, using functional magnetic resonance imaging, deep/slow breathing was shown to increase cortical and subcortical activity (Critchley et al. 2015), which may help to support improved physical and mental health (Laborde et al. 2022). Pursed-lips breathing typically involves nasal inspiration and prolonged expiration directed through lips that have a “puckered” or “pursed” appearance (see Fig. 3). Independent of other breathing strategies, expiration through pursed lips may increase expiratory resistance at the mouth, evoking a small positive end-expiratory pressure of ~5 cm H₂O (van der Schans et al. 1997). This can help ameliorate dynamic airway compression and expiratory flow limitation in patients with COPD by functionally “stenting” the airways (Marciniuk et al. 2011; Nguyen and Duong 2021). Diaphragmatic breathing comprises many of the aforementioned techniques but with preferential activation of the diaphragm during inspiration. From a practical standpoint, diaphragmatic




Fig. 3 Primary outcomes from the literature on systematized breathing interventions. ↑ evidence of increase; ↓ evidence of decrease, ↔ evidence of no change, FEV_1 forced expiratory volume in 1 s, *COPD* chronic obstructive pulmonary disease

breathing is achieved by inspiring with minimal movement of the chest and more pronounced outward abdominal displacement (Cahalin et al. 2002).

Literature

The literature on breathing interventions employs inconsistent nomenclature, with terms often being used interchangeably. This makes it difficult to discern the efficacy of each technique. Accordingly, we have grouped the literature on breathing interventions and discussed their collective outcomes. The exceptions are pursed-lips breathing (see “[Pursed-lips breathing \(PLB\)](#)”) and the Buteyko Breathing Technique (see “[The Buteyko Breathing Technique](#)”), both of which have enough independent research to warrant their own discussions. Most of the literature on breathing interventions has focused on the management of respiratory symptoms in asthma and COPD. Regarding the former, the Cochrane database published a meta-analysis and separate systematic review on breathing exercises in asthmatic adults and children, respectively, with disparate findings. In adults, yoga (including pranayama), breathing retraining, the Buteyko Breathing

Technique, the Papworth method, and deep diaphragmatic breathing, all improved quality of life, symptoms and complaints due to hyperventilation (defined in the study as “over breathing”), and lung function (forced expiratory volume in 1 s; [FEV_1]). However, studies were characterized by poor methodologies and a very low-to-moderate quality of evidence (Santino et al. 2020). The review in children found insufficient data to support the use of breathing techniques for asthma management, owing primarily to a low number of studies ($n = 3$, 112 participants) (Macêdo et al. 2016). Another issue impeding interpretation of the pediatric data is that studies generally combined breathing exercises with a comprehensive package of care, thereby precluding any evaluation of breathing exercises alone (Macêdo et al. 2016). As such, due to a low number of studies, limited reporting of data, and variations in reported outcomes, no firm conclusions can be drawn regarding the efficacy of breathing interventions for asthma management. More well-controlled, high-quality studies are needed.

There is a much larger body of work evaluating breathing interventions for improving respiratory symptoms, lung function, and exercise capacity in COPD. In general,

long-term breathing interventions including deep breathing with or without preferential activation of the diaphragm, pursed-lips breathing, yoga, singing, and breathing gymnastics, all appear to improve functional exercise performance (mainly 6-min walk test; [6MWT]) (Hamasaki 2020; Lu et al. 2020; Yang et al. 2022), quality of life (St. George's Respiratory Questionnaire) (Marotta et al. 2020) and stress and anxiety (Hamasaki 2020). Several studies also show that pursed-lips breathing, with or without preferential activation of the diaphragm, improves pulmonary function (i.e., forced vital capacity [FVC] and FEV₁) (Hamasaki 2020; Lu et al. 2020; see Yang et al. 2022).

The literature on breathing interventions for improving dyspnea in COPD is less consistent. A review of 13 RCTs ($n = 998$ patients) found that home-based breathing exercises (diaphragmatic breathing, yoga breathing, breathing gymnastics, and singing) improved resting FEV₁, 6MWT distance, and ratings of dyspnea (modified Medical Research Council dyspnea scale and St George's Respiratory Questionnaire) across the range of disease severity (Lu et al. 2020). By contrast, a Cochrane review of 16 studies ($n = 1233$ patients) showed that 15 weeks of breathing retraining (pursed-lips breathing, diaphragmatic breathing, ventilation feedback training, or yoga breathing, both supervised and unsupervised) improved 6MWT distance in COPD but had no consistent effects on dyspnea at rest or health-related quality of life (Holland et al. 2012). Another systematic review and meta-analysis of 19 studies ($n = 745$ patients) reported that respiratory frequency was significantly reduced at rest and during exercise following a period of dedicated pursed-lips breathing, ventilatory feedback and exercise, diaphragmatic breathing, or combined techniques that lasted between one day and 24 weeks—nevertheless, breathing interventions did not improve ratings of dyspnea relative to controls (Ubolnuar et al. 2019). The reason for the discrepancy in dyspnea-related findings is unclear; however, Holland et al. (2012) and Ubolnuar et al. (2019) included studies performed in various environments (such as the laboratory and during pulmonary rehabilitation programs in outpatient settings), whereas the review by Lu et al. (2020) focused exclusively on home-based interventions.

Pursed-lips breathing (PLB)

This particular technique has received a great deal of attention as a standalone therapy in patients with COPD owing to its effects on dyspnea and exercise tolerance. The main benefits include reduced respiratory frequency, increased (improved) inspiratory and total respiratory time, and increased tidal volume (Ubolnuar et al. 2019). A bout of PLB has been shown to reduce resting CO₂ retention and increase arterial oxygen tension and arterial oxyhemoglobin saturation in advanced but stable COPD (Thoman et al.

1966; Breslin 1992; Marciniuk et al. 2011). The primary mechanism by which PLB exerts its effects is by increasing intraluminal airway pressure during expiration which tends to prevent the airway compression that would otherwise occur as intrapleural pressure increases. In turn, PLB is likely to ameliorate air trapping. Pursed-lips breathing also reduces expiratory reserve volume and lengthens the diaphragm (thereby improving its tension-generating capacity during inspiration) (Spahija et al. 2005); subsequently, increased arterial oxyhemoglobin saturation may be the result of a more complete, mechanically-efficient respiratory cycle. Pursed-lips breathing has been used by COPD patients during exercise. Generally, favorable outcomes have been observed on 6MWT (Bhatt et al. 2013), perhaps mediated by reduced dynamic lung hyperinflation (Cabral et al. 2015), increased arterial oxyhemoglobin saturation (Cabral et al. 2015), and possible protection against diaphragm fatigue (Breslin 1992). Notwithstanding, improvements in exercise capacity with PLB are not a universal finding (Garrod et al. 2005).

Several studies in patients with COPD have tried to distinguish the benefits of PLB from other respiratory interventions or relaxation techniques. Pivotal research by Tiep et al. (1986) showed that an acute (15-min) bout of PLB evoked greater increases in SpO₂ at rest compared with general relaxation techniques. Others have found that 12 weeks of daily practice in PLB was more effective at reducing exertional dyspnea, and increasing 6MWT performance, than expiratory muscle training or a control group that received an educational pamphlet but no intervention (Niell et al. 2007). In another study, a 12-week PLB intervention increased FEV₁ and maximal inspiratory pressures by a greater magnitude than diaphragmatic breathing or a no-intervention control (Jansang et al. 2016). One study assessed the effects of an acute bout of diaphragmatic breathing with or without PLB on COPD patients during upright, seated rest (Mendes et al. 2019). Although both interventions increased ribcage and abdominal volumes (measured via respiratory inductive plethysmography), increased arterial oxyhemoglobin saturation, and decreased respiratory frequency, there were greater reductions in respiratory frequency and longer expiratory times with combined diaphragmatic and pursed-lips breathing. Conversely, neither intervention reduced dyspnea (Medical Research Council Scale) or end-expiratory chest volume. Thus, the combination of breath control (with preferential activation of the diaphragm) and prolonged expiration through pursed lips may provide additive benefits that are distinct from other breathing techniques. Indeed, as an effective means of managing dyspnea in COPD, PLB has been advocated by the Canadian Thoracic Society in their clinical practice guidelines (Marciniuk et al. 2011) and by

the American Thoracic Society in their patient education materials (Lareau et al. 2020).

The Buteyko Breathing Technique

This (predominantly) commercial breathing regimen was conceived in the 1950s by Dr Konstantin Pavlovic Buteyko. In its modern form, Buteyko is an amalgam of several breathing techniques which emphasize nasal breathing and periods of breath-hold (referred to as “control pause”). In general, the research on Buteyko is favorable, particularly with respect to asthma management, showing improved quality of life scores (Burgess et al. 2011; Santino et al. 2020). Nevertheless, Buteyko breathing does not appear to be superior to other chronic breathing interventions like yoga, deep/slow breathing, PLB, or diaphragmatic breathing. In addition, several issues cloud the interpretation of the Buteyko literature, potentially undermining its validity.

First, Buteyko is usually administered as a comprehensive package of care that comprises breathing retraining, education, and nutritional advice, making it difficult to discern the isolated benefits of the respiratory intervention (Bruton and Lewith 2005). Second, proponents of Buteyko often extend the claims beyond those supported by the scientific literature. For instance, a major premise of the technique is that breath-hold time predicts alveolar CO₂ according to a patented mathematical formula—a claim that has been empirically disproven (Courtney and Cohen 2008). The Buteyko Breathing Technique also advocates mouth taping as a means of obligating nasal breathing during sleep. However, a randomized, crossover study in patients with symptomatic asthma showed that mouth taping had no effect on asthma control (Cooper et al. 2009). Some proponents even suggest, without evidence, that Buteyko can treat diseases and symptoms (including diabetes, attention-deficit hyperactive disorder, and dental health)—claims which undermine the scientific legitimacy of the intervention. It is also worth noting that most clinical studies on Buteyko have assessed outcomes in response to physiotherapy programs that tend to focus on the more conventional, evidence-based aspects of the technique (e.g., deep breathing through the nose).

Another approach of some Buteyko proponents is to associate the technique with unrelated, or tenuously related, research. For example, studies have identified a high prevalence of allergic rhinitis in children with ADHD (Brawley et al. 2004). By promoting nasal breathing, Buteyko advocates imply that the technique can reduce the risk of developing ADHD. Lastly, several commercial incarnations of Buteyko promote long breath holds (> 25 s) which may be unsuitable for certain groups (e.g., COPD patients). Accordingly, while the more conventional aspects of Buteyko (nasal inspiration, deep/slow breathing, and breath training) may have benefits for respiratory function, it is important

that health and exercise professionals remain wary of claims that are lacking plausibility, currently unproven, and potentially dangerous.

Additional considerations and conclusions on respiratory interventions

The literature on breathing techniques for patients with respiratory disease is generally positive. Still, there are subtle nuances in the data that should be highlighted prior to interpretation. For example, research suggests that respiratory physiotherapy using breathing training has the potential to improve inspiratory and expiratory muscle strength following upper abdominal surgery (Grams et al. 2012), yet this has questionable utility following general abdominal surgery (Pasquina et al. 2006). Physicians and other healthcare professionals must also be conscious of instances where breathing training may be less favorable or even harmful to their patients. Respiratory physiotherapy seems to have limited benefits on lung volume or mortality risk following lung resection (Larsen et al. 2020), and PLB specifically has been shown to increase metabolic demands in patients with interstitial lung disease (Parisien-La Salle et al. 2019). This brings into question the efficacy of PLB for restrictive disorders. Certain breathing techniques may also worsen respiratory symptoms; e.g., several studies show that diaphragmatic breathing may exacerbate dyspnea in patients with severe COPD (Hamasaki 2020), perhaps due to negative effects on the work of breathing and the muscle's mechanical efficiency (Gosselink et al. 2012). Anecdotally, the technical demands of preferentially activating the diaphragm during inspiration may render such interventions unsuitable for respiratory patients.

Evidence summary and recommendations

Primary outcomes from the literature on breathing interventions are summarized in Fig. 3. Breathing interventions such as deep breathing and pursed-lips breathing may elicit favorable changes in tidal volume, respiratory frequency, respiratory time, and arterial oxygen saturation in patients with COPD, particularly those with severe or very severe disease. Long-term breathing retraining strategies may improve lung function (mainly lung volumes and capacities), exercise performance, respiratory symptoms, and quality of life in respiratory patients. Pursed-lips breathing, in particular, is an important standalone therapy that should be considered as an adjunct to exercise training and pharmaceutical interventions in pulmonary rehabilitation programs. Breathing interventions should emphasize a deep and slow nasal inspiration followed by a slow and prolonged expiration through pursed lips. Because of possible negative outcomes in patients, breathing interventions should be delivered by

experienced therapists with a comprehensive understanding of the benefits and risks of each technique. Patients must also be managed on a case-by-case basis. The benefits of deep/slow breathing in healthy subjects (individuals with 'normal' pulmonary function) are likely centered on increased parasympathetic activity and the associated physiological perturbations (e.g., increased heart rate variability, reduced resting heart rate, reduced resting blood pressure) which may, in turn, support improved mental health.

Respiratory muscle training

Premise and plausibility

The healthy respiratory system has typically been considered “overbuilt” for the ventilatory demands placed upon it during strenuous exercise. More recently, however, studies have revealed several respiratory constraints that may impede exercise performance in healthy subjects, particularly those who are endurance-trained, and in certain patients with cardiorespiratory disease (Dempsey et al. 2020). When breathing frequency increases during exercise, there is a consequent increase in the resistive loads placed upon the inspiratory and expiratory muscles. In patients with COPD, the resistive loads are exacerbated due to narrowing of peripheral airways. In addition, at lung volumes above relaxation volume, where lung and chest wall compliance are reduced, the inspiratory muscles must overcome increased elastic recoil forces. This increase in elastic loading occurs when tidal volume increases with exercise, and especially in the presence of dynamic lung hyperinflation (i.e., increase in end-expiratory lung volume) consequent to the dynamic compression of airways during forced expiration. Airway narrowing and loss of elastic recoil in COPD give rise to static lung hyperinflation, which further increases the elastic loading on the inspiratory muscles. In severe COPD, incomplete expiration and inward recoil of the lungs and chest wall result in progressive air trapping and increased alveolar pressure at the end of expiration (i.e., intrinsic positive end-expiratory pressure, PEEPi). To initiate inspiratory airflow, the inspiratory muscles must generate a negative pressure equal in magnitude to PEEPi, thereby imposing a threshold load on the inspiratory muscles. When the lung is acutely inflated, the pressure-generating capacity of the diaphragm is impaired because the muscle is shortened. At high lung volumes, the pressure-generating capacity of the diaphragm may be further reduced by an increased radius of muscle curvature. Lung inflation also impairs the pressure-generating capacity of the inspiratory intercostal muscles (external intercostals and parasternal intercostals); in contrast to the diaphragm, however, this impairment has been ascribed to changes in the orientation and motion of the ribs (De Troyer and Wilson 2009). In COPD, reductions

in the pressure-generating capacity of respiratory muscles may also result from disease-induced changes in respiratory muscle morphology.

Increased loading of the respiratory muscles and/or decreased capacity of the respiratory muscles for pressure generation have direct functional consequences. For instance, an increase in the respiratory muscle load/capacity ratio contributes significantly to the subjective experience of breathing discomfort (i.e., dyspnea) (McConnell and Romer 2004a) and predisposes the respiratory muscles to fatigue. Regarding the latter, decreases in the contractile function of inspiratory or expiratory muscles have been noted following intense, whole-body exercise in healthy young adults (Johnson et al. 1993; Taylor et al. 2006; Tiller et al. 2017) and in select patients with COPD (Hopkinson et al. 2010; Bachasson et al. 2013). This exercise-induced decrease in respiratory muscle capacity further increases the intensity of dyspnea. Moreover, the metabolite accumulation associated with fatiguing respiratory muscle work can elicit a sympathetically-mediated vasoconstrictor response in locomotor muscles. This ‘respiratory muscle metaboreflex’ may decrease locomotor muscle blood flow in favor of an increase in blood flow to the respiratory muscles, thereby increasing the fatigability of limb locomotor muscles and reducing central motor output via feedback effects (Sheel et al. 2018). Excessive loading of the respiratory muscles and activation of the respiratory muscle metaboreflex may partly contribute to the early development of limb muscle fatigue in patients with COPD (Amann et al. 2010).

Respiratory muscle training (RMT) was developed on the premise that enhancing the pressure-generating capacity of respiratory muscles would increase fatigue resistance and/or mechanical efficiency of the respiratory muscles during exercise. Such changes would be expected to prevent or delay the respiratory muscle metaboreflex, thereby improving O₂ delivery to working limbs and reducing the intensity of perceived limb discomfort (see Fig. 4). An increase in the pressure-generating capacity of the respiratory muscles with targeted training would also be expected to reduce the intensity of perceived dyspnea. In health, RMT has been shown to improve the static and dynamic function of respiratory muscles (Romer and McConnell 2003), attenuate fatigability of respiratory (Verges et al. 2007, 2009) and locomotor muscles (McConnell and Lomax 2006), blunt the respiratory muscle metaboreflex (Witt et al. 2007), and attenuate sensations of respiratory and limb discomfort (McConnell and Romer 2004a). Research pertaining to the influence of RMT on whole-body exercise performance is somewhat contradictory (see Literature).

The three most common approaches to RMT involve flow-resistive loading (high pressure, low flow), pressure-threshold loading (high pressure, moderate flow), and isocapnic voluntary hyperpnea (low pressure, high flow).




Fig. 4 Primary outcomes from the literature on respiratory muscle training. One of the putative mechanisms underpinning the effects of respiratory muscle training on exercise tolerance and performance is

a possible ‘blunting’ of the respiratory muscle metaboreflex. ↑ evidence of increase, ↓ evidence of decrease, ↔ evidence of no change. *COPD* chronic obstructive pulmonary disease

Devices that impose a resistive or threshold load elicit improvements predominately in respiratory muscle strength, whereas isocapnic voluntary hyperpnea elicits improvements predominantly in respiratory muscle endurance (see McConnell and Romer 2004b for review). More recently, a tapered flow-resistive loading device has been developed to produce a variable load that matches the pressure–volume relationship of inspiratory muscles (Langer et al. 2013). A recent development in the RMT literature pertains to external loading of the respiratory muscles during exercise (in-task). So-called “functional” RMT typically involves flow-resistive loading via facemask (Porcari et al. 2016)¹ or nasal restriction (Arnedillo et al. 2020; Gonzalez-Montesinos et al. 2021). Although functional RMT is an attractive proposition due to its specificity of application, the additional loads

imposed on the respiratory muscles, and hence the potential training stimulus, are difficult to quantify. From a practical standpoint, functional RMT may limit the physiological stimulus that can be obtained by applying RMT and exercise independently (Faghy et al. 2021).

¹ Applied external resistors are intentionally designed to elicit high resistive loads during exercise; thus, they impose considerably greater loads than low-resistance face coverings (e.g., cloth and surgical masks) that might be used for personal protection from airborne pathogens. Indeed, the negative physiological effects of protective face masks have been shown to be negligible when used during physical activity in healthy individuals (Hopkins et al. 2021).

Literature

In 1976, Leith and Bradley (Leith and Bradley 1976) showed that the respiratory muscles of healthy individuals could be trained to increase strength or endurance. Later research sought to evaluate the efficacy of RMT with respect to whole-body exercise performance in healthy individuals and in patients with respiratory disease. Unfortunately, many of the early studies were hampered by methodological shortcomings, including small sample sizes, absence of sham-control groups, unbalanced baseline characteristics, inadequate training intensities, and inappropriate outcome measures (McConnell and Romer 2004b). As such, the ergogenic effect of RMT has been the subject of much debate (e.g., McConnell 2012; Patel et al. 2012). A systematic review and meta-analysis of 46 studies on the effects of RMT in healthy subjects revealed an improvement in endurance performance as assessed using fixed-intensity tests, simulated time-trials, and intermittent incremental tests (Illi et al. 2012). The analysis also showed that resistive/threshold and hyperpnea training did not differ in their effects, that combined inspiratory/expiratory strength training tended to be superior to either intervention alone, and that the greatest improvements with RMT occurred in less-fit subjects and in sports of longer duration (Illi et al. 2012). Another systematic review and meta-analysis, this time on responses in athletes, showed a positive effect of RMT on respiratory muscle function and sport performance outcomes (HajBhanbari et al. 2013). Although the report also noted comparable benefits of RMT for “elite” and “recreational athletes”, the authors classified training status by whether the subject’s $\dot{V}O_2$ max was above or below the minimum, pre-determined requirements for being considered an “athlete”, but without specifically defining “elite”. Thus, the question as to whether training status mediates the efficacy of RMT remains unresolved. More recent studies have shown improvements in repeated-sprint performance (e.g., shorter recovery between sprints or increased number of repetitions) as well as reduced effort perceptions and markers of metabolic stress after resistive RMT (Lorca-Santiago et al. 2020). Collectively, the data show an ergogenic effect of RMT on endurance and repeated-sprint performance in healthy individuals.

The efficacy of RMT in patients with COPD has been studied extensively. Although whole-body exercise training is a crucial component of pulmonary rehabilitation (Casaburi 2008), exercise training does not appear to increase the pressure-generating capacity of the respiratory muscles in this population. Consequently, there has been a great deal of interest in the potential for RMT to increase the capacity of the respiratory muscles and alleviate symptoms. Learned societies, including the American College of Chest Physicians/American Association of Cardiovascular

and Pulmonary Rehabilitation (Ries et al. 2007) and the European Respiratory Society/American Thoracic Society (Spruit et al. 2013), have recommended RMT for patients who, despite optimal medical therapy, exhibit dyspnea and reduced respiratory muscle strength. Several systematic reviews and meta-analyses have shown that RMT, when applied as a standalone intervention with controlled training loads in patients with COPD, improves respiratory muscle strength and endurance, exercise capacity, dyspnea, and health-related quality of life (e.g., Gosselink et al. 2011). Adding RMT to a whole-body exercise training program in COPD has been shown to have no additive effects on exercise performance or quality of life, suggesting that RMT may only be effective as a standalone treatment in the absence of other interventions (e.g., Gosselink et al. 2011). This notion is corroborated by a recent systematic review and meta-analysis which showed that inspiratory pressure-threshold training in patients with COPD increased inspiratory muscle strength, functional exercise performance, and dyspnea during activities of daily living, but with no additional effect on the intensity of exertional dyspnea when used as an adjunct to pulmonary rehabilitation (Beaumont et al. 2018a). Notwithstanding the limitations of meta-analyses (e.g., poor quality of included studies, study heterogeneity, publication bias), recent large-scale RCTs on the effects of RMT in patients with COPD have confirmed that improvements in inspiratory muscle function after adjunctive RMT do not translate to additional improvements in functional exercise capacity, dyspnea, or quality of life (Beaumont et al. 2018b; Schultz et al. 2018; Charususin et al. 2018). In patients with inspiratory muscle weakness, however, adjunctive RMT during a whole-body exercise training intervention elicited a significant increase in endurance cycling time and a significant reduction in dyspnea intensity at iso-time during the cycling test compared to sham-training (Charususin et al. 2018).

Additional applications and population subgroups

While most studies have investigated the influence of RMT on exercise outcomes in healthy individuals (athletes and non-athletes) and patients with COPD, RMT may also have application in other settings where the loads imposed on the respiratory muscles are elevated or the capacity to generate force is reduced. For instance, RMT has been considered in the context of environmental and occupational settings (e.g., altitude and load carriage) (Faghy and Brown 2016; Chambault et al. 2021) and in the context of natural aging (Seixas et al. 2020; Manifold et al. 2021). In older adults, exertional dyspnea is consistently elevated during submaximal exercise owing to an increased ventilatory demand (Jensen et al. 2009). Conceivably, RMT might improve exertional dyspnea in older adults through a reduction in the load/capacity ratio

of respiratory muscles. In addition to the increase in ventilatory demand, aging is accompanied by a decline in respiratory muscle function which, through a reduction in postural control, has been shown to correlate with impairments in balance performance (Rodrigues et al. 2020). In turn, RMT has been shown to improve balance performance through an increase in the neuromuscular activity of postural muscles (Ferraro et al. 2019, 2020, 2022; Tounsi et al. 2021).

Recent evidence indicates that females have smaller airways than males, and subsequently exhibit greater flow-resistive work of breathing (Peters et al. 2021). There also appears to be a combined influence of age and biological sex on respiratory mechanics which contributes in part to the increased perception of exertional dyspnea noted in older women (Molgat-Seon et al. 2018). Thus, RMT could be an effective intervention to enhance the overall exercise response in young and older women. Other groups with imbalances in the load/capacity ratio of respiratory muscles, and which might therefore benefit from RMT, include: exercise-induced laryngeal obstruction (Sandnes et al. 2022), obstructive sleep apnea (Torres-Castro et al. 2022), cystic fibrosis (Stanford et al. 2020), interstitial lung disease (Zaki et al. 2022), stroke (Fabero-Garrido et al. 2022), hypertension (Craighead et al. 2022), chronic heart failure (Azambuja et al. 2020), pulmonary hypertension (Tran et al. 2021), neurological disorders (He et al. 2021), spinal cord injury (Woods et al. 2022), pre-operative surgery (Dsouza et al. 2021), weaning from mechanical ventilation (Worraphan et al. 2020), ventilator-induced diaphragm dysfunction in the recovery phase (Ahmed et al. 2019), and COVID-19 (e.g., risk reduction, ICU, recovery, and long-COVID) (McNarry et al. 2022). Evidence of the efficacy of RMT in these groups requires further prospective study.

Evidence summary and recommendations

Primary outcomes from the literature on RMT are summarized in Fig. 4. If applied with the appropriate frequency, intensity, and duration, RMT can improve specific aspects of respiratory muscle function (e.g., strength and endurance). There is convincing evidence of an ergogenic effect of RMT in healthy individuals (athletes and non-athletes). As a standalone therapy, RMT confers multiple benefits for select patients with COPD. However, the effect of adding RMT to a general exercise program in COPD (e.g., within pulmonary rehabilitation) appears limited. It is conceivable that RMT may be useful for patients with respiratory muscle weakness or those with pre-existing comorbidities who are unable to participate in whole-body exercise training. Further RCTs are needed to ascertain which patients and groups are likely to benefit from RMT.

Canned oxygen

Premise and plausibility

Commercial canned oxygen (intended for non-medical use) is a can of hyperoxic gas (~95% O₂) equipped with a mask or inhaler cap. The suggested protocol for use differs among manufacturers but typically involves several inhalations, repeated 8–10 times, periodically throughout the day or as needed. Some vendors recommend their product for use immediately before physical activity and/or sporting competition. The ergogenic claims include improved reaction time, “improved breathing” during exposure to heat and pollution, and improved sports performance by delaying the onset of fatigue and improving O₂ availability for oxidative metabolism. Some brands combine eucalyptus and other oils with the gas mixture which they claim can “relax the nervous system, relax the muscles, and relieve stress”. Despite the extensive claims and widespread and costly prescription of so-called “short burst oxygen therapy” for respiratory patients (e.g., COPD), there is no clear mechanism for the purported physiological benefit. Moreover, in healthy individuals, hemoglobin remains nearly completely saturated with O₂ at rest, and exercise-induced arterial O₂ desaturation (i.e., hypoxemia) rarely occurs in healthy (untrained) individuals at sea-level. Consequently, it seems implausible that acute exposure to concentrated O₂ (i.e., several breaths) could influence respiratory outcomes or exercise performance.

Literature

The focus of this section is on commercially available canned oxygen and “short burst oxygen” rather than physician-prescribed supplemental oxygen therapy. A systematic review on the efficacy of short-burst oxygen to improve breathlessness, exercise capacity, arterial oxygen saturation, and ventilatory variables in patients with COPD concluded that its widespread prescription was not evidence-based (O’Neill et al. 2006). Due to a lack of peer-reviewed studies on commercial canned oxygen in particular, most vendors cite clinical literature that is tenuously related (e.g., studies on hyperbaric oxygen therapy or prolonged inhalation of medically certified gas mixtures). Thus, the references provided by manufacturers do not support the claims. One manufacturer published an online press release that was designed to have the appearance of a scientific journal article, presumably in an effort to feign scientific legitimacy. On the rare occasion that relevant journal articles were obtained through commercial websites, they were of very low quality and exhibited a high risk of bias. It is worth noting that although gaseous supplemental oxygen (delivered by inhalation) is not prohibited by the World Anti-Doping

Agency (WADA 2022), some sports authorities prohibit its use. Athletes should therefore be cognizant of the rules and regulations regarding O₂ therapy that govern their sport.

Evidence summary and recommendations

The proposed benefit of acute inhalation of canned oxygen has low plausibility and there is no valid evidence of beneficial effects.

Nutritional interventions

Premise and plausibility

Nutrition is a modifiable factor that influences the development and progression of many non-communicable diseases (Cena and Calder 2020; Dominguez et al. 2021). Some nutrients have immunomodulatory, anti-inflammatory, and/or antioxidant effects (Kau et al. 2011; Venter et al. 2020; Gozzi-Silva et al. 2021). Such nutrients may therefore influence respiratory health and disease risk/progression in conditions underpinned by airway and/or systemic inflammation (Berthon and Wood 2015; Hosseini et al. 2017; Parvizian et al. 2020; Heloneida de Araújo Morais et al. 2021). In addition, supplementation with certain nutrients may provide prophylactic and/or therapeutic benefits for some respiratory patients.

In terms of therapeutic effects on respiratory health, the bulk of literature focuses on vitamin D, antioxidants (most commonly vitamin C), omega-3 polyunsaturated fatty acids (n-3 PUFAs), probiotics, and prebiotics. The wealth of literature precludes a detailed discussion of the complex and diverse mechanisms underpinning each nutrient and their independent effects on respiratory health. Instead, the following summary focuses on the purported antioxidant and immunomodulatory effects and whether they translate to clinically meaningful outcomes.

Literature on vitamin D₃

Circulating concentrations of 25(OH)D—a form of vitamin D produced in the liver from hydroxylation of vitamin D₃—were found to be inversely associated with the incidence of upper- and/or lower-respiratory tract infection (RTI) (Pham et al. 2019). The incidence of RTI in the general population peaks in the winter (Ginde et al. 2009) when vitamin D deficiency is most common owing to low skin exposure to sunlight ultraviolet B radiation (Farrokhyar et al. 2014; Cashman et al. 2016). Indeed, insufficient circulating concentrations of vitamin D (baseline serum 25(OH)D < 50 nmol·L⁻¹) have been observed in military personnel (Harrison et al. 2021), athletes (Farrokhyar et al. 2014), and healthy controls (Cashman et al. 2016).

Vitamin D sufficiency can be achieved via oral vitamin D₃ supplementation (Carswell et al. 2018; Harrison et al. 2021) and safe exposure to sunlight or simulated sunlight of the appropriate wavelength. However, evidence of prophylactic and/or therapeutic effects of vitamin D₃ supplementation is confounded by heterogeneity across trials (Jolliffe et al. 2021), with effect-modifiers including dosing regimen and duration, participant age, baseline 25(OH)D, and geographic location (Martineau et al. 2017; Vlieg-Boerstra et al. 2021; Jolliffe et al. 2021; Cho et al. 2022). A recent meta-analysis of 43 RCTs ($n = 48,488$ mixed-health cohort of children and adults) revealed a modest but overall decreased risk of acute respiratory infection with daily vitamin D₃ supplementation of 400–1000 IU (Jolliffe et al. 2021). A recent RCT reported no change in the incidence of upper-respiratory tract infection following 12 weeks of vitamin D₃ supplementation (1000 IU d⁻¹ for the first four weeks and 400 IU d⁻¹ for eight weeks), but found decreased peak severity and duration of illness (Harrison et al. 2021). Accordingly, irrespective of whether vitamin D₃ supplementation influences the incidence of upper-RTI, it may still attenuate its severity and/or duration.

There is a high prevalence of vitamin D deficiency in patients with asthma (Bener et al. 2014) and COPD (Janssens et al. 2011). In these patients, higher vitamin D concentrations are associated with lower risk, severity, and exacerbation of the primary disease (Gupta et al. 2011; Zhu et al. 2016; Liu et al. 2019). Vitamin D supplementation in these groups has thus been studied for its prophylactic and therapeutic effects. There is insufficient evidence that the prophylactic use of vitamin D₃ can prevent asthma in children (Yepes-Nuñez et al. 2018; Luo et al. 2022). In addition, the association between vitamin D status and adult-onset asthma is unclear (Mai et al. 2012; Cheng et al. 2014; Confino-Cohen et al. 2014; Cherrie et al. 2017; Manousaki et al. 2017). The therapeutic effects of vitamin D₃ supplementation in children and adults with pre-existing asthma are also equivocal (Jolliffe et al. 2021; Chen et al. 2021). For instance, a recent systematic review (Nitzan et al. 2022) and an independent meta-analysis (Kumar et al. 2021) both concluded that vitamin D₃ supplementation did not affect lung function, asthma control, or exacerbation rates in children. Although these studies were not performed in children with pre-existing vitamin D deficiency, findings generally concur with recent RCTs in children with 25(OH)D concentration < 50 nmol L⁻¹ (Jat et al. 2021) and < 75 nmol L⁻¹ (Forno et al. 2020; Han et al. 2021). By contrast, in asthmatic adults with low vitamin D₃ concentration, a meta-analysis of three small trials ($n = 92$) revealed some protection of vitamin D₃ supplementation against exacerbations (Jolliffe et al. 2017). One RCT also showed improved asthma control in 25(OH)D-deficient adults who were supplemented with a weekly dose of 16,000 IU (Andújar-Espinosa et al. 2021).

Based on the aforementioned evidence, vitamin D₃ supplementation does not generally improve lung function in COPD patients (Lehouck et al. 2012; Sluyter et al. 2017; Chen et al. 2019; Foumani et al. 2019), but it may confer improvements in FEV₁ in current or former smokers with 25(OH)D < 50 nmol L⁻¹ (Sluyter et al. 2017). Data from one meta-analysis of four RCTs (*n* = 560) indicate that vitamin D₃ supplementation reduces exacerbation rates in vitamin D-deficient patients (Jolliffe et al. 2019). Thus, for COPD patients who are hospitalized for exacerbation, the Global Initiative for Chronic Obstructive Lung Disease recommends vitamin D screening and subsequent supplementation in those found to be deficient (Global Initiative for Chronic Obstructive Lung Disease 2022).

Literature on vitamin C

In healthy populations, the efficacy of regular vitamin C (ascorbic acid) supplementation on upper-RTI incidence depends on individual physical stress levels and associated immune perturbations. Broadly speaking, the data show no benefit of chronic supplementation. A meta-analysis of 24 trials (> 10,000 participants) showed no effect of moderate- or high-dose vitamin C on the incidence of upper-RTI (Hemilä and Chalker 2013). Accordingly, chronic vitamin C supplementation is not justified in normal (vitamin C-replete) populations (Hemilä and Chalker 2013; Gómez et al. 2018). The short-term, therapeutic effects of vitamin C are less conclusive (Hemilä and Chalker 2013), but several reviews report that supplementation shortened the duration of upper-RTI symptoms by ~ 8–18% (Hemilä and Chalker 2013; Abioye et al. 2021). It may therefore be practical to initiate short-term vitamin C supplementation within 24 h of symptom onset.

There is also evidence that acute supplementation may benefit individuals undergoing periods of extreme physical stress. An analysis of five trials comprising marathon runners, skiers, and soldiers, found that vitamin C supplementation reduced symptoms of the common cold by ~ 50% following "severe physical exercise" (Hemilä and Chalker 2013). Further to moderating virally-mediated respiratory symptoms, the prophylactic effects of vitamin C supplementation in some athletes may result from an attenuation of EIB (Tecklenburg et al. 2007; Hemilä 2013). Thus, when the risk of infection in athletes is elevated due to a high training/competitive load (Ruuskanen et al. 2022) or extensive travel (Walsh 2019), vitamin C supplementation (0.25–1.0 g d⁻¹) may reduce the severity and/or duration of upper-RTIs (Walsh 2019; Cerullo et al. 2020). An important caveat is that chronic, high-dose (~ 1 g d⁻¹) vitamin C supplementation may blunt certain training-induced skeletal muscle adaptations (Mason et al. 2020), and is therefore discouraged.

Lastly, there is some evidence that vitamin C may help ameliorate asthma symptoms (Allen et al. 2009; Berthon and Wood 2015). An analysis of three small trials (*n* = 40) in asthmatics found that vitamin C supplementation, in various dosing regimens (1.5 g d⁻¹ for two weeks; 2 g ingested 1 h before exercise; 0.5 g ingested 1.5 h before exercise), attenuated the post-exercise fall in FEV₁ by 48% (Hemilä 2013). Notwithstanding, there is insufficient evidence to make decisive recommendations regarding vitamin C supplementation for asthma management, and more RCTs with larger samples are needed.

Literature on omega-3 (n-3) poly-unsaturated fatty acids (PUFAs)

The most abundant PUFA in the Western diet is linoleic acid which is converted to arachidonic acid—a precursor for pro-inflammatory and bronchoconstrictive signaling. By contrast, omega-3 (n-3) PUFAs, including eicosapentaenoic acid (EPA) and docosahexaenoic acid (DHA), derived primarily from fatty fish, may have anti-inflammatory effects. Specifically, EPA inhibits arachidonic acid, blunts pro-inflammatory signaling, and acts as a precursor for pro-resolving mediators with anti-inflammatory properties (Brannan et al. 2015). Similarly, DHA has been shown to modify gene expression and signaling pathways related to inflammatory mediators (Calder 2010). Dietary supplementation with EPA and DHA has therefore been explored as an adjunct therapy in certain respiratory conditions (Thien et al. 2002; Yang et al. 2013; Stoodley et al. 2019).

Studies show that supplementation with high dose n-3 PUFAs for several weeks mitigates EIB (Mickleborough et al. 2003, 2006; Tecklenburg-Lund et al. 2010; Mickleborough and Lindley 2014; Kumar et al. 2016). Yet, because high dose n-3 PUFA is expensive and may cause gastrointestinal complaints, it is worth noting that both high dose (6.2 g/d) and moderate dose (3.1 g/d) n-3 PUFA appear to exert similar effects on provocation-induced decreases in FEV₁ (Williams et al. 2017).

In COPD, studies with n-3 PUFA supplementation show equivocal results. A meta-analysis of eight RCTs found that supplementation increased body mass, increased low-density lipoproteins, and reduced IL-6, but did not influence lung function or quality of life (Yu et al. 2021). These results should be interpreted cautiously because, depending on disease severity and other comorbidities, weight gain may be beneficial for some COPD patients and harmful for others. Moreover, some RCTs provide limited data regarding individual doses of EPA and DHA. For example, an observational cohort study of > 120,000 US women and men initially showed that greater consumption of fish (> 4 servings per week) was associated with lower risk of newly diagnosed COPD. But subsequent analysis revealed that COPD risk

was unrelated to total n-3 PUFA intake (Varraso et al. 2015). To date, only one observational cohort study in moderate-to-severe COPD has shown that high dietary n-3 PUFA reduces risk of severe exacerbations, decreases the number of respiratory symptoms, improves health-related quality of life, and reduces overall morbidity (Lemoine et al. 2020). The same study showed the opposite effects of high dietary n-6 PUFA (linoleic acid) (Lemoine et al. 2020). These data speak to the importance of distinguishing n-3 from n-6 PUFA in supplementation interventions, and the importance of the dietary n-3/n-6 PUFA ratio in respiratory health. More well-controlled RCTs on n-3 PUFA supplementation in current and former smokers with COPD are warranted.

Literature on probiotics, prebiotics, and synbiotics

The microbial profile and gut microbiome have a substantial influence on health and disease (Clemente et al. 2012) and systemic immune function (Roberfroid et al. 2010). Immune function is particularly important for respiratory health and disease, and the “gut-lung axis” represents a promising (non-pharmacological) therapeutic target (Marsland et al. 2015). Beneficial changes in the gut microbiota can be achieved through dietary supplementation with probiotics (live microorganisms that confer a health benefit on the host when administered in adequate amounts) (Hill et al. 2014), prebiotics (substrates that are selectively utilized by host microorganisms, conferring a health benefit) (Gibson et al. 2017), and/or synbiotics (a combination of pro- and prebiotics).

In terms of the gut microbiota and its effects on respiratory health in subjects without respiratory disease, the largest body of evidence relates to upper-RTIs, and the data are largely favorable. For example, a 2015 Cochrane review of 10 trials found that probiotics reduced the incidence of upper-RTI relative to placebo (Hao et al. 2015). Meta-analyses show similar findings in healthy infants, children, and adults after supplementation with probiotics (six studies, $n = 1682$) (Rashidi et al. 2021) and synbiotics (four RCTs, $n = 883$) (Chan et al. 2020). Probiotics may also decrease upper-RTI risk in active individuals and athletes (Cox et al. 2010; West et al. 2011, 2014; Haywood et al. 2014; Strasser et al. 2016).

There is also preliminary data indicating that the gut-lung axis may be a suitable target for managing asthma and related conditions. Prebiotics, probiotics, and synbiotics each reduced airway inflammation and disease severity in rodent models of allergic asthma (Sagar et al. 2014; Verheijden et al. 2015, 2016). Furthermore, a small-scale, double-blind, placebo-controlled, randomized trial showed potential benefits of prebiotics in adults with EIB (Williams et al. 2016). Specifically, prebiotics reduced serum markers of airway inflammation at baseline and completely

abolished the 29% provocation-induced increase in TNF- α (a pro-inflammatory cytokine). Lastly, eight weeks supplementation with probiotics decreased asthma exacerbations in children when compared to placebo (Drago et al. 2022). Although more RCTs in humans are warranted, the pre-clinical rodent data and preliminary human in-vivo studies show potential benefits of prebiotics and/or probiotics as potential adjunct therapies to support respiratory health.

Evidence summary and recommendations

Primary outcomes from the literature on nutritional interventions are summarized in Fig. 5. The effects of chronic vitamin D₃ supplementation on the prevalence/severity of upper-RTI are inconsistent. When supplemented prophylactically, there is no evidence of benefit in asthma management. In asthmatics and COPD patients with pre-existing deficiency, vitamin D₃ supplementation may confer therapeutic benefits. Long-term, daily supplementation of vitamin C (ascorbic acid) provides little-to-no benefit in those who are vitamin C-replete, but may reduce the severity and/or duration of the common cold and symptoms of general RTI when supplemented acutely at symptom onset ($0.25\text{--}1.0\text{ g d}^{-1}$), especially in individuals undergoing periods of extreme physical stress. Nevertheless, there is insufficient evidence supporting vitamin C supplementation for asthma management. Several weeks of n-3 PUFA supplementation reduces the severity of EIB. However, similar data in COPD are equivocal, with only one observational cohort study showing benefits in terms of reduced risk of exacerbation, respiratory symptoms, and overall morbidity. Daily probiotics and/or prebiotics reduce the incidence of upper-RTI more than placebo in adults, children, active individuals, and athletes. Prebiotics, probiotics, and synbiotics may also reduce airway inflammation and disease severity in rodent models of allergic asthma, with preliminary evidence showing benefits in adults with EIB.

Inhaled L-menthol

Premise and plausibility

L-menthol is a cyclic alcohol derived from the oils of various species of *Mentha* (mints) that have been used as medicinal plants for millennia. There is evidence that inhaled or ingested L-menthol triggers a cooling sensation by stimulating sensory nerve endings in the nasal vestibule and mucosa that convey nasal sensation (Aldren and Tolley 1993; Eccles 2003). Because L-menthol has a significant positive effect on the sensation of nasal airflow, some menthol vendors claim that the oil can decongest the upper airways (e.g., during colds and allergies), enhance nasal flow, and improve airway patency. L-menthol is widely available in lozenges, nasal




Fig. 5 Primary outcomes from the literature on nutritional interventions. ↑ evidence of increase; ↓ evidence of decrease; ↔ evidence of no change. RTI respiratory tract infection; EIB exercise-induced bronchoconstriction

sprays, vapor rubs, inhalers, cough syrups, mouthwashes, as a scent in aromatherapy oils, and as a flavoring in cigarettes and e-cigarettes. However, the plausibility of L-menthol to improve respiratory function is low because it does not possess the amine group that would be expected of a substance with vasodilator or bronchodilator properties, nor does it have a chemical structure similar to nasal decongestants (Eccles et al. 1988; Eccles 1994). Accordingly, any benefit of L-menthol is likely to be indirect—mediated by cooling sensations that stimulate the nasal trigeminal nerve thereby creating the cognitive illusion of improved inspiratory flow (Kanezaki et al. 2021).

Literature

Articles were excluded if L-menthol was not inhaled, dissolved and nebulized, ingested orally as a lozenge, rinsed/swilled in the mouth, if the effects of L-menthol could not be distinguished from other substances that were co-administered, or if the article did not assess respiratory function. It is well-established that healthy adults (free from respiratory disorders and the common cold) experience increased sensations of nasal airflow and/or nasal patency after inhaling L-menthol (Eccles et al. 1988; Pereira et al. 2013). Two randomized, placebo-controlled trials showed that inhaling

L-menthol reduced sensations of respiratory discomfort during flow-resistive and elastic loading at rest (Nishino et al. 1997) and inspiratory resistive loading during exercise (Kanezaki and Ebihara 2017). Studies in individuals with the common cold also found that a menthol-containing lozenge evoked marked improvements in sensations of nasal airflow and decongestion (Eccles et al. 1990; Eccles et al. 1990). Yet, subjective changes in respiratory sensations are not reflected in objective changes in breathing pattern (i.e., respiratory frequency, tidal volume, or inspiratory flow), minute ventilation, or spirometric indices of lung function (Nishino et al. 1997; Kanezaki and Ebihara 2017). Case in point, Köteles et al. (2018) showed that nebulized menthol-containing peppermint, rosemary, or eucalyptus oil, inhaled over 15 min, had no effect on FVC, FEV₁/FVC, or peak expiratory flow (PEF), despite improving the perceptions of spirometric outcomes. Similarly, the only study to assess upper-airway resistance using rhinometry confirmed there was no effect of menthol on nasal/upper-airway resistance, respiratory frequency, or minute ventilation in healthy adults at rest (Pereira et al. 2013).

In obstructive respiratory disorders, the data tend to follow a similar pattern. During inspiratory resistive loading in patients with mild-to-severe COPD, L-menthol significantly improved subjective measures (i.e., physical and mental “breathing effort”, air hunger, breathing discomfort, and anxiety and fear) relative to a non-L-menthol control, but did not influence objective measures (i.e., breathing pattern, respiratory duty cycle, and inspiratory muscle activity) (Kanezaki et al. 2020). Others have observed no difference between nebulized menthol and placebo with respect to changes in FVC or FEV₁ in mild asthmatics (Tamaoki et al. 1995). A randomized, double-blind trial in patients with chronic cough found that, in response to a capsaicin provocation test, inhalation of 1 mL nebulized L-menthol (0.5% and 1% concentration) improved peak inspiratory flow relative to placebo, whereas only high-dose L-menthol (1%) attenuated the reduction in forced inspiratory flow at 50% of vital capacity (FIF₅₀) and increased the cough threshold (Millqvist et al. 2013). A randomized, single-blinded, placebo-controlled crossover trial in patients with chronic bronchitis, found that “aromatics” containing L-menthol significantly improved mucous clearance when compared to petroleum jelly, but without subsequent changes in lung function (Hasani et al. 2003). Lastly, although studies have generally failed to observe any direct effect of L-menthol on physiological variables during exercise, there is a possible indirect effect of L-menthol on exercise performance in the heat owing to changes in the sensation of oropharyngeal temperature versus placebo (Mündel and Jones 2010).

Evidence summary and recommendations

Primary outcomes from the literature on inhaled L-menthol are summarized in Fig. 6. By stimulating sensory nerve endings in the nasal vestibule and mucosa (e.g., the trigeminal nerve), inhaled L-menthol can augment sensations of nasal airflow, improve respiratory sensations in both healthy subjects and patients, and potentially relieve dyspnea in COPD. Improved respiratory sensations may translate to improved exercise performance in the heat. There is some evidence that high-concentration inhaled L-menthol may increase the cough threshold in patients with chronic cough. Nevertheless, L-menthol does not have vasodilator or bronchodilator properties, and there is little-to-no convincing evidence that L-menthol has direct functional benefits on spirometry-derived variables in any population.

Conclusions

The health and wellness industry is characterized by, and in many cases depends on, lax consumer regulations on products and services. As a result, interventions are often sold on insufficient evidence, baseless claims, and pseudoscience (Tiller et al. 2022). Not only is there a growing disparity between the substance of commercial claims and the supporting scientific evidence, thereby violating Laplace’s principle that “Extraordinary claims require extraordinary evidence”, but the legitimate (plausible) and illegitimate (implausible) claims for these interventions are often conflated, obscuring the translation of science to practice. This is a particular problem in the field of respiratory physiology and medicine.

This review is intended as an evidence-based guide to help health and exercise professionals distinguish science from pseudoscience in commercial respiratory interventions and make informed decisions that optimize patient/client outcomes. In summarizing the recommendations, there are several caveats that should be noted. First, the products/strategies selected for inclusion were commercial interventions (i.e., not controlled drugs or products regulated by the FDA as “medical devices”). The list was delimited to those interventions most prevalent in the health and wellness industry that were coupled to the most conspicuous claims, and there may be prominent, mainstream interventions that were not included.

A second caveat is that the recommendations herein are based on data from controlled laboratory-based studies. The statistical analyses typically used in such studies have allowed researchers to reject, or fail to reject, the null hypotheses, and subsequently discuss the existence of effects or lack thereof. Yet, such an approach is dichotomous by design, providing little room for nuanced interpretation of differences, potentially




Fig. 6 Primary outcomes from the literature on inhaled L-menthol. ↑ evidence of increase; ↓ evidence of decrease; ↔ evidence of no change. FEV₁ forced expiratory volume in 1 s, FVC forced vital capacity, PEF peak expiratory flow, COPD chronic obstructive pulmonary disease

overlooking practical or clinical significance. For example, some studies in exercise rehabilitation have been shown to yield non-significant between-group differences despite moderate-to-large effects that would be deemed meaningful in practice (Zemková 2014). Interventions with moderate-to-large effects, despite lack of statistical significance, may be especially important for high-performance athletes for whom the margins of success are extremely small. The opposite may also be true (i.e., statistical tests may yield highly significant outcomes with trivial effects). To improve external validity in exercise-based studies, researchers have been encouraged to include confidence intervals and/or effect sizes in their statistical analyses as measures of “practical significance” (Knudson 2009). This might aid in the interpretation of both “statistically significant” and “practically meaningful” outcomes.

In this comprehensive review of literature and expert consensus, it was determined that: (1) there is good quality data supporting subjective/perceptual (but not objective) benefits of both nasal dilators and L-menthol; (2) there is some evidence that nasally-derived nitric oxide may benefit critically ill patients but not healthy subjects; (3) there is good evidence that systematized breathing interventions (particularly pursed-lips breathing) can improve exercise

performance, respiratory symptoms, and quality of life in COPD and asthma; (4) there is good evidence that respiratory muscle training can improve exercise performance in healthy subjects, respiratory symptoms in some patient populations (e.g., COPD), and benefit COPD patients with respiratory muscle weakness or pre-existing comorbidities, the latter of which may preclude them from whole-body exercise training; (5) there is some evidence that nutritional interventions, including vitamin D and vitamin C, may benefit respiratory health in individuals with pre-existing nutrient deficiencies and in healthy individuals during times of compromised immune function associated with increased physical stress, and interesting but inconsistent evidence of benefits of polyunsaturated fatty acids and pre/probiotics/synbiotics; and (6) no evidence that canned oxygen is beneficial for any clinical outcome.

For the interventions aforementioned, we advocate for greater vigilance in determining prior plausibility and evidence for efficacy. We also hope to inspire similar expert reviews that scrutinize interventions in other facets of the commercial health and wellness industry.

Author contributions CRI and NBT: conceived the manuscript. All authors drafted, edited, and approved the final version.

Funding No funding was received to develop this manuscript. Nicholas B. Tiller is funded by a postdoctoral fellowship from the Tobacco-Related Disease Research Program (TRDRP; award no. T31FT1692). Camilla R. Illidi is funded by an Accelerate postdoctoral fellowship from the Mathematics of Information Technology and Complex System (MITACS award no. IT28775).

Data availability The datasets generated during and/or analyzed during the current study are available from the corresponding author on reasonable request.

Declarations

Conflict of interest The authors declare no conflicts of interest and do not have any financial disclosures.

References

- Abioye AI, Bromage S, Fawzi W (2021) Effect of micronutrient supplements on influenza and other respiratory tract infections among adults: a systematic review and meta-analysis. *BMJ Glob Health* 6:e003176. <https://doi.org/10.1136/bmjgh-2020-003176>
- Adams CM, Peiffer JJ (2017) Neither internal nor external nasal dilation improves cycling 20-km time trial performance. *J Sci Med Sport* 20:415–419. <https://doi.org/10.1016/j.jsams.2016.08.018>
- Ahmed S, Martin AD, Smith BK (2019) Inspiratory muscle training in patients with prolonged mechanical ventilation: narrative review. *Cardiopulm Phys Ther J* 30:44–50. <https://doi.org/10.1097/CPT.0000000000000092>
- Aldren C, Tolley NS (1993) Further studies on nasal sensation of airflow. *Rhinology* 29:49–55
- Allen S, Britton JR, Leonardi-Bee JA (2009) Association between antioxidant vitamins and asthma outcome measures: systematic review and meta-analysis. *Thorax* 64:610–619. <https://doi.org/10.1136/THX.2008.101469>
- Amann M, Regan MS, Kobitary M et al (2010) Impact of pulmonary system limitations on locomotor muscle fatigue in patients with COPD. *Am J Physiol Regul Integr Comp Physiol* 299:R314–324. <https://doi.org/10.1152/ajpregu.00183.2010.-We>
- Amaro ACS, Duarte FHG, Jallad RS et al (2012) The use of nasal dilator strips as a placebo for trials evaluating continuous positive airway pressure. *Clinics* 67:469–474. [https://doi.org/10.6061/clinics/2012\(05\)11](https://doi.org/10.6061/clinics/2012(05)11)
- Anderson SD, Kippelen P (2012) Assessment and prevention of exercise-induced bronchoconstriction. *Br J Sports Med* 46:391–396. <https://doi.org/10.1136/bjsports-2011-090810>
- Andújar-Espinosa R, Salinero-González L, Illán-Gómez F et al (2021) Effect of vitamin D supplementation on asthma control in patients with vitamin D deficiency: the ACVID randomised clinical trial. *Thorax* 76:126–133. <https://doi.org/10.1136/THORAXJNL-2019-213936>
- Arnedillo A, Gonzalez-Montesinos JL, Fernandez-Santos JR et al (2020) Effects of a rehabilitation programme with a nasal inspiratory restriction device on exercise capacity and quality of life in COPD. *Int J Environ Res Public Health* 17:3669. <https://doi.org/10.3390/IJERPH17103669>
- Azambuja ADCM, De Oliveira LZ, Sbruzzi G (2020) Inspiratory muscle training in patients with heart failure: what is new? Systematic review and meta-analysis. *Phys Ther* 100:2099–2109. <https://doi.org/10.1093/PTJ/PZAA171>
- Bachasson D, Wuyam B, Pepin JL et al (2013) Quadriceps and respiratory muscle fatigue following high-intensity cycling in COPD patients. *PLoS One* 8:e83432. <https://doi.org/10.1371/journal.pone.0083432>
- Bahammam AA, Tate R, Manfreda J, Kryger MH (1999) Upper airway resistance syndrome: effect of nasal dilation, sleep stage, and sleep position. *Sleep* 22:592–598
- Banzett RB, Lansing RW, Binks AP (2021) Air hunger: a primal sensation and a primary element of dyspnea. *Compr Physiol* 11:1449–1483. <https://doi.org/10.1002/CPHY.C200001>
- Barnes CS, Alexis NE, Bernstein JA et al (2013) Climate change and our environment: the effect of respiratory and allergic disease. *J Allergy Clin Immunol Pract* 1:137–141. <https://doi.org/10.1016/j.jaip.2012.07.002.Climate>
- Beaumont M, Forget P, Couturaud F, Reyckler G (2018a) Effects of inspiratory muscle training in COPD patients: a systematic review and meta-analysis. *Clin Respir J* 12:2178–2188. <https://doi.org/10.1111/CRJ.12905>
- Beaumont M, Mialon P, Le Ber C et al (2018b) Effects of inspiratory muscle training on dyspnoea in severe COPD patients during pulmonary rehabilitation: controlled randomised trial. *Eur Respir J* 51:1701107. <https://doi.org/10.1183/13993003.01107-2017>
- Becquemin MH, Swift DL, Bouchikhi A et al (1991) Particle deposition and resistance in the noses of adults and children. *Eur Respir J* 4:694–702
- Bener A, Ehlayel MS, Bener HZ, Hamid Q (2014) The impact of Vitamin D deficiency on asthma, allergic rhinitis and wheezing in children: an emerging public health problem. *J Fam Community Med* 21:154–161. <https://doi.org/10.4103/2230-8229.142967>
- Bennett WD, Zeman KL, Jarabek AM (2003) Nasal contribution to breathing with exercise: effect of race and gender. *J Appl Physiol* 95:497–503. <https://doi.org/10.1152/jappphysiol.00718.2002>
- Berthon BS, Wood LG (2015) Nutrition and respiratory health – feature review. *Nutrients* 7:1618–1643. <https://doi.org/10.3390/nu7031618>
- Bhatt SP, Luqman-Arafath TK, Gupta AK et al (2013) Volitional pursed lips breathing in patients with stable chronic obstructive pulmonary disease improves exercise capacity. *Chron Respir Dis* 10:5–10. <https://doi.org/10.1177/1479972312464244>
- Bishop CA, Johnson SM, Wall MB et al (2016) Magnetic resonance imaging reveals the complementary effects of decongestant and Breathe Right Nasal Strips on internal nasal anatomy. *Laryngoscope* 126:2205–2211. <https://doi.org/10.1002/lary.25906>
- Boggs GW, Ward JR, Stavrianeas S (2008) The external nasal dilator: style over function? *J Strength Cond Res* 22:269–275. <https://doi.org/10.1519/JSC.0b013e31815f903e>
- Brannan JD, Bood J, Alkhabaz A et al (2015) The effect of omega-3 fatty acids on bronchial hyperresponsiveness, sputum eosinophilia, and mast cell mediators in asthma. *Chest* 147:397–405. <https://doi.org/10.1378/CHEST.14-1214>
- Brawley A, Silverman B, Kearney S et al (2004) Allergic rhinitis in children with attention-deficit/hyperactivity disorder. *Ann Allergy Asthma Immunol* 92:663–667. [https://doi.org/10.1016/S1081-1206\(10\)61434-2](https://doi.org/10.1016/S1081-1206(10)61434-2)
- Breslin EH (1992) The pattern of respiratory muscle recruitment during pursed-lip breathing. *Chest* 101:75–78. <https://doi.org/10.1378/chest.101.1.75>
- Bruton A, Lewith GT (2005) The Buteyko breathing technique for asthma: a review. *Complement Ther Med* 13:41–46. <https://doi.org/10.1016/J.CTIM.2005.01.003>
- Burgess J, Ekanayake B, Lowe A et al (2011) Systematic review of the effectiveness of breathing retraining in asthma management. *Expert Rev Respir Med* 5:789–807. <https://doi.org/10.1586/ERS.11.69>

- Cabral LF, D'Elia TDC, Marins DDS et al (2015) Pursed lip breathing improves exercise tolerance in COPD: a randomized crossover study. *Eur J Phys Rehabil Med* 51:79–88
- Cahalin LP, Braga M, Matsuo Y, Hernandez ED (2002) Efficacy of diaphragmatic breathing in persons with chronic obstructive pulmonary disease: a review of the literature. *J Cardpulm Rehabil* 22:7–21. <https://doi.org/10.1097/00008483-200201000-00002>
- Calder PC (2010) Omega-3 fatty acids and inflammatory processes. *Nutrients* 2:355–374. <https://doi.org/10.3390/NU2030355>
- Camacho M, Malu OO, Kram YA et al (2016) Nasal dilators (Breathe Right strips and NoZovent) for snoring and OSA: a systematic review and meta-analysis. *Pulm Med* 2016:4841310. <https://doi.org/10.1155/2016/4841310>
- Carswell AT, Oliver SJ, Wentz LM et al (2018) Influence of vitamin D supplementation by sunlight or oral D3 on exercise performance. *Med Sci Sports Exerc* 50:2555–2564. <https://doi.org/10.1249/MSS.0000000000001721>
- Casaburi R (2008) A brief history of pulmonary rehabilitation. *Respir Care* 53:1185–1189
- Cashman KD, Dowling KG, Skrabáková Z et al (2016) Vitamin D deficiency in Europe: pandemic? *Am J Clin Nutr* 103:1033–1077. <https://doi.org/10.3945/ajcn.115.120873>
- Cena H, Calder PC (2020) Defining a healthy diet: Evidence for the role of contemporary dietary patterns in health and disease. *Nutrients* 12:334. <https://doi.org/10.3390/nu12020334>
- Cerullo G, Negro M, Parimbelli M et al (2020) The long history of vitamin C: from prevention of the common cold to potential aid in the treatment of COVID-19. *Front Immunol* 11:574029. <https://doi.org/10.3389/FIMMU.2020.574029>
- Chambault J, Grand G, Kayser B (2021) Sex-specific effects of respiratory muscle endurance training on cycling time trial performance in normoxia and hypoxia. *Front Physiol* 12:1–8. <https://doi.org/10.3389/fphys.2021.700620>
- Chan CKY, Tao J, Chan OS et al (2020) Preventing respiratory tract infections by synbiotic interventions: a systematic review and meta-analysis of randomized controlled trials. *Adv Nutr* 11:979–988. <https://doi.org/10.1093/ADVANCES/NMAA003>
- Charususin N, Gosselink R, Decramer M et al (2018) Randomised controlled trial of adjunctive inspiratory muscle training for patients with COPD. *Thorax* 73:942–950. <https://doi.org/10.1136/THORAXJNL-2017-211417>
- Chen FY, Xiao M, Ling B et al (2019) Vitamin D does not improve lung function decline in COPD: a meta-analysis. *Eur Rev Med Pharmacol Sci* 23:8637–8644. https://doi.org/10.26355/EUR-REV_201910_19181
- Chen Z, Peng C, Mei J et al (2021) Vitamin D can safely reduce asthma exacerbations among corticosteroid-using children and adults with asthma: a systematic review and meta-analysis of randomized controlled trials. *Nutr Res* 92:49–61. <https://doi.org/10.1016/J.NUTRES.2021.05.010>
- Cheng HM, Kim S, Park GH et al (2014) Low vitamin D levels are associated with atopic dermatitis, but not allergic rhinitis, asthma, or IgE sensitization, in the adult Korean population. *J Allergy Clin Immunol* 133:1048–1055. <https://doi.org/10.1016/J.JACI.2013.10.055>
- Cherrie MPC, Sarran C, Osborne NJ (2017) Supplementary materials: association between serum 25-hydroxyvitamin D levels and the prevalence of adult-onset asthma. *Int J Environ Res Public Health* 15:1103. <https://doi.org/10.3390/ijerph15061103>
- Cho H, Myung SK, Cho HE (2022) Efficacy of vitamin D supplements in treatment of acute respiratory infection: a meta-analysis for randomized controlled trials. *Nutrients* 14:1144. <https://doi.org/10.3390/nu14061144>
- Clemente JC, Ursell LK, Parfrey LW, Knight R (2012) The impact of the gut microbiota on human health: an integrative view. *Cell* 148:1258–1270. <https://doi.org/10.1016/J.CELL.2012.01.035>
- Confino-Cohen R, Brufman I, Goldberg A, Feldman BS (2014) Vitamin D, asthma prevalence and asthma exacerbations: a large adult population-based study. *Allergy* 69:1673–1680. <https://doi.org/10.1111/ALL.12508>
- Cooper S, Osborne J, Harrison T, Tattersfield A (2009) Effect of mouth taping at night on asthma control – a randomised single-blind crossover study. *Respir Med* 103:813–819. <https://doi.org/10.1016/J.RMED.2009.02.003>
- Courtney R, Cohen M (2008) Investigating the claims of Konstantin Buteyko, M.D., Ph.D.: the relationship of breath holding time to end tidal CO2 and other proposed measures of dysfunctional breathing. *J Altern Complement Med* 14:115–123. <https://doi.org/10.1089/ACM.2007.7204>
- Cox AJ, Pyne DB, Saunders PU, Fricker PA (2010) Oral administration of the probiotic *Lactobacillus fermentum* VRI-003 and mucosal immunity in endurance athletes. *Br J Sports Med* 44:222–226. <https://doi.org/10.1136/BJSM.2007.044628>
- Craighead DH, Tavoian D, Freeberg KA et al (2022) A multi-trial, retrospective analysis of the antihypertensive effects of high-resistance, low-volume inspiratory muscle strength training. *J Appl Physiol* (Bethesda Md: 1985). <https://doi.org/10.1152/JAPPLPHYSIOL.00425.2022>
- Crespo AS, Hallberg J, Lundberg JO et al (2010) Nasal nitric oxide and regulation of human pulmonary blood flow in the upright position. *J Appl Physiol* 108:181–188. <https://doi.org/10.1152/jappphysiol.00285.2009>
- Critchley HD, Nicotra A, Chiesa PA et al (2015) Slow breathing and hypoxic challenge: cardiorespiratory consequences and their central neural substrates. *PLoS One* 10:e0127082. <https://doi.org/10.1371/journal.pone.0127082>
- Dallam GM, McClaran SR, Cox DG, Foust CP (2018) Effect of nasal versus oral breathing on Vo2max and physiological economy in recreational runners following an extended period spent using nasally restricted breathing. *Int J Kinesiol Sports Sci* 6:22–29. <https://doi.org/10.7575/aiac.ijkss.v.6n.2p.22>
- de Araújo H, Morais A, de Souza AJ, Kelly da Silva-Maia J et al (2021) Nutritional status, diet and viral respiratory infections: perspectives for severe acute respiratory syndrome coronavirus 2. *Br J Nutr* 125:851–862. <https://doi.org/10.1017/S0007114520003311>
- De Troyer A, Wilson TA (2009) Effect of acute inflation on the mechanics of the inspiratory muscles. *J Appl Physiol* 107:315–323. <https://doi.org/10.1152/JAPPLPHYSIOL.91472.2008>
- Dempsey JA, La GA, Hull JH (2020) Is the healthy respiratory system built just right, overbuilt, or underbuilt to meet the demands imposed by exercise? *J Appl Physiol* 129:1235–1256. <https://doi.org/10.1152/JAPPLPHYSIOL.00444.2020>
- Di Somma EM, West SN, Wheatley JR, Amis TC (1999) Nasal dilator strips increase maximum inspiratory flow via nasal wall stabilization. *Laryngoscope* 109:780–784. <https://doi.org/10.1097/00005537-199905000-00018>
- Dinardi RR, Ribeiro De Andrade C, Da Cunha IC (2013) Evaluation of the effectiveness of the external nasal dilator strip in adolescent athletes: a randomized trial. *Int J Pediatr Otorhinolaryngol* 77:1500–1505. <https://doi.org/10.1016/j.ijporl.2013.06.018>
- Dinardi RR, De Andrade CR, Da Cunha Ibiapina C (2014) External nasal dilators: definition, background, and current uses. *Int J Gen Med* 7:491–504. <https://doi.org/10.2147/IJGM.S67543>
- Dinardi RR, de Andrade CR, da Cunha IC (2017) Effect of the external nasal dilator on adolescent athletes with and without allergic rhinitis. *Int J Pediatr Otorhinolaryngol* 97:127–134. <https://doi.org/10.1016/j.ijporl.2017.04.008>

- Dinardi RR, Santos Ferreira CH, Santos Silveira G et al (2021) Does the external nasal dilator strip help in sports activity? A systematic review and meta-analysis. *Eur Arch Otorhinolaryngol* 278:1307–1320. <https://doi.org/10.1007/s00405-020-06202-5>
- Dominguez LJ, Di Bella G, Veronese N, Barbagallo M (2021) Impact of Mediterranean diet on chronic non-communicable diseases and longevity. *Nutrients* 13:2028. <https://doi.org/10.3390/nu13062028>
- Drago L, Cioffi L, Giuliano M et al (2022) A post hoc analysis on the effects of a probiotic mixture on asthma exacerbation frequency in schoolchildren. *ERJ Open Res* 8:00020–02022. <https://doi.org/10.1183/23120541.00020-2022>
- Dsouza FV, Amaravadi SK, Samuel SR et al (2021) Effectiveness of expiratory muscle training on respiratory muscle strength in patients undergoing cardiac surgeries: a systematic review with meta-analysis. *Ann Rehabil Med* 45:264–273. <https://doi.org/10.5535/ARM.21027>
- Eccles R (1994) Menthol and related cooling compounds. *J Pharm Pharmacol* 46:618–630. <https://doi.org/10.1111/J.2042-7158.1994.TB03871.X>
- Eccles R (2003) Menthol: effects on nasal sensation of airflow and the drive to breathe. *Curr Allergy Asthma Rep* 3:210–214. <https://doi.org/10.1007/S11882-003-0041-6>
- Eccles R, Griffiths DH, Newton CG, Tolley NS (1988) The effects of menthol isomers on nasal sensation of airflow. *Clin Otolaryngol Allied Sci* 13:25–29. <https://doi.org/10.1111/j.1365-2273.1988.tb00277.x>
- Eccles R, Jawad MS, Morris S (1990) The effects of oral administration of (–)-menthol on nasal resistance to airflow and nasal sensation of airflow in subjects suffering from nasal congestion associated with the common cold. *J Pharm Pharmacol* 42:652–654. <https://doi.org/10.1111/j.2042-7158.1990.tb06625.x>
- Eccles R, Morris S, Jawad MSM (1990) The effects of menthol on reaction time and nasal sensation of airflow in subjects suffering from the common cold. *Clin Otolaryngol Allied Sci* 15:39–42. <https://doi.org/10.1111/j.1365-2273.1990.tb00430.x>
- Ehteshami-Afshar S, FitzGerald JM, Doyle-Waters MM, Sadatsafavi M (2016) The global economic burden of asthma and chronic obstructive pulmonary disease. *Int J Tuberc Lung Dis* 20:11–23. <https://doi.org/10.5588/ijtld.15.0472>
- Elizondo S, Ayala-Sanchez D, Choi W, et al (2019) Physiological effects of supplemental concentrated oxygen in Vo2 kinetics during steady-state exercise
- Fabero-Garrido R, del Corral T, Angulo-Díaz-Parreño S et al (2022) Respiratory muscle training improves exercise tolerance and respiratory muscle function/structure post-stroke at short term: a systematic review and meta-analysis. *Ann Phys Rehabil Med*. <https://doi.org/10.1016/J.REHAB.2021.101596>
- Faghy MA, Brown PI (2016) Training the inspiratory muscles improves running performance when carrying a 25 kg thoracic load in a backpack. *Eur J Sport Sci* 16:585–594. <https://doi.org/10.1080/17461391.2015.1071878>
- Faghy MA, Brown PI, Davis NM et al (2021) A flow resistive inspiratory muscle training mask worn during high-intensity interval training does not improve 5 km running time-trial performance. *Eur J Appl Physiol* 121:183–191. <https://doi.org/10.1007/S00421-020-04505-3>
- Farrokhyar F, Tabasinejad R, Dao D et al (2014) Prevalence of vitamin D inadequacy in athletes: a systematic-review and meta-analysis. *Sports Med* 45:365–378. <https://doi.org/10.1007/S40279-014-0267-6>
- Ferraro FV, Gavin JP, Wainwright T, McConnell A (2019) The effects of 8 weeks of inspiratory muscle training on the balance of healthy older adults: a randomized, double-blind, placebo-controlled study. *Physiol Rep* 7:e14076. <https://doi.org/10.14814/PHY2.14076>
- Ferraro FV, Gavin JP, Wainwright TW, McConnell AK (2020) Comparison of balance changes after inspiratory muscle or Otago exercise training. *PLoS One* 15:1–16. <https://doi.org/10.1371/journal.pone.0227379>
- Ferraro FV, Gavin JP, Wainwright TW, McConnell AK (2022) Association between inspiratory muscle function and balance ability in older people: a pooled data analysis before and after inspiratory muscle training. *J Ageing Phys Act* 30:421–433. <https://doi.org/10.1123/JAPA.2020-0507>
- Forno E, Bacharier LB, Phipatanakul W et al (2020) Effect of vitamin D3 supplementation on severe asthma exacerbations in children with asthma and low vitamin D levels: the VDKA randomized clinical trial. *J Am Med Assoc* 324:752–760. <https://doi.org/10.1001/JAMA.2020.12384>
- Foumani AA, Mehrdad M, Jafarizhad A et al (2019) Impact of vitamin D on spirometry findings and quality of life in patients with chronic obstructive pulmonary disease: a randomized, double-blind, placebo-controlled clinical trial. *Int J Chron Obstruct Pulmon Dis* 14:1495–1501. <https://doi.org/10.2147/COPD.S207400>
- Garrod R, Dallimore K, Cook J et al (2005) An evaluation of the acute impact of pursed lips breathing on walking distance in non-spontaneous pursed lips breathing chronic obstructive pulmonary disease patients. *Chron Respir Dis* 2:67–72. <https://doi.org/10.1191/1479972305cd0680a>
- Gelardi M, Porro G, Sterlicchio B et al (2019) Internal and external nasal dilator in patients who snore: a comparison in clinical practice. *Acta Biomed* 90:10–14. <https://doi.org/10.23750/abm.v90i2-S.8096>
- Gerlach H, Rossaint R, Pappert D, Falke KJ (1993) Time-course and dose-response of nitric oxide inhalation for systemic oxygenation and pulmonary hypertension in patients with adult respiratory distress syndrome. *Eur J Clin Invest* 23:499–502. <https://doi.org/10.1111/j.1365-2362.1993.tb00797.x>
- Gibson GR, Hutkins R, Sanders ME et al (2017) Expert consensus document: the International Scientific Association for Probiotics and Prebiotics (ISAPP) consensus statement on the definition and scope of prebiotics. *Nat Rev Gastroenterol Hepatol* 14:491–502. <https://doi.org/10.1038/NRGASTRO.2017.75>
- Ginde AA, Mansbach JM, Camargo CA (2009) Association between serum 25-hydroxyvitamin D level and upper respiratory tract infection in the third national health and nutrition examination survey. *Arch Intern Med* 169:384–390. <https://doi.org/10.1001/archinternmed.2008.560>
- Global Initiative for Chronic Obstructive Lung Disease Inc (2022) Global Strategy for the Diagnosis, management, and Prevention of Chronic Obstructive Pulmonary Disease
- Gómez E, Quidel S, Bravo-Soto G, Ortigoza Á (2018) Does vitamin C prevent the common cold? *Medwave* 18:e7235. <https://doi.org/10.5867/MEDWAVE.2018.04.7236>
- Gonzalez-Montesinos JL, Fernandez-Santos JR, Vaz-Pardal C et al (2021) Chronic effects of a training program using a nasal inspiratory restriction device on elite cyclists. *Int J Environ Res Public Health* 18:1–13. <https://doi.org/10.3390/IJERPH18020777>
- Gosselink R, De Vos J, Van Den Heuvel SP et al (2011) Impact of inspiratory muscle training in patients with COPD: what is the evidence? *Eur Respir J* 37:416–425. <https://doi.org/10.1183/09031936.00031810>
- Gosselink R, Wagenaar RC, Rijswijk H et al (2012) Diaphragmatic breathing reduces efficiency of breathing in patients with chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 151:1136–1142. <https://doi.org/10.1164/AJRCM.151.4.7697243>
- Gozzi-Silva SC, Teixeira FME, da Silva Duarte AJ et al (2021) Immunomodulatory role of nutrients: how can pulmonary dysfunctions

- improve? *Front Nutr* 8:674258. <https://doi.org/10.3389/FNUT.2021.674258>
- Grams ST, Ono LM, Noronha MA et al (2012) Breathing exercises in upper abdominal surgery: a systematic review and meta-analysis. *Rev Bras Fisioter* 16:345–353. <https://doi.org/10.1590/S1413-35552012005000052>
- Griffin JW, Hunter G, Ferguson D, Sillers MJ (1997) Physiologic effects of an external nasal dilator. *Laryngoscope* 107:1235–1238. <https://doi.org/10.1097/00005537-199709000-00014>
- Gupta A, Sjoukes A, Richards D et al (2011) Relationship between serum vitamin D, disease severity, and airway remodeling in children with asthma. *Am J Respir Crit Care Med* 184:1342–1349. <https://doi.org/10.1164/rccm.201107-1239OC>
- Gustafsson LE, Leone AM, Persson MG et al (1991) Endogenous nitric oxide is present in the exhaled air of rabbits, guinea pigs and humans. *Biochem Biophys Res Commun* 181:852–857. [https://doi.org/10.1016/0006-291X\(91\)91268-H](https://doi.org/10.1016/0006-291X(91)91268-H)
- HajBhanbari B, Yamabayashi C, Buna TR et al (2013) Effects of respiratory muscle training on performance in athletes: a systematic review with meta-analyses. *J Strength Cond Res* 27:1643–1663. <https://doi.org/10.1519/JSC.0b013e318269f73f>
- Hamasaki H (2020) Effects of diaphragmatic breathing on health: a narrative review. *Medicines* 7:1–19. <https://doi.org/10.3390/medicines7100065>
- Han YY, Forno E, Bacharier LB et al (2021) Vitamin D supplementation, lung function and asthma control in children with asthma and low vitamin D levels. *Eur Respir J* 58:2100989. <https://doi.org/10.1183/13993003.00989-2021>
- Hao Q, Dong BR, Wu T (2015) Probiotics for preventing acute upper respiratory tract infections. *Cochrane Database Syst Rev*. <https://doi.org/10.1002/14651858.CD006895.PUB3>
- Harrison SE, Oliver SJ, Kashi DS et al (2021) Influence of vitamin D supplementation by simulated sunlight or oral D3 on respiratory infection during military training. *Med Sci Sports Exerc* 53:1505–1516. <https://doi.org/10.1249/MSS.0000000000002604>
- Hasani A, Pavia D, Toms N et al (2003) Effect of aromatics on lung mucociliary clearance in patients with chronic airways obstruction. *J Altern Complement Med* 9:243–249. <https://doi.org/10.1089/10755530360623356>
- Haywood BA, Black KE, Baker D et al (2014) Probiotic supplementation reduces the duration and incidence of infections but not severity in elite rugby union players. *J Sci Med Sport* 17:356–360. <https://doi.org/10.1016/J.JSAMS.2013.08.004>
- He Y, Zhao C, Liu Y (2021) Effects of respiratory muscle training on cough function in neurological disorders: a systematic review with meta-analysis. *NeuroRehabilitation* 48:441–449. <https://doi.org/10.3233/NRE-210017>
- Hemilä H (2013) Vitamin C may alleviate exercise-induced bronchoconstriction: a meta-analysis. *BMJ Open* 3:e002416. <https://doi.org/10.1136/BMJOPEN-2012-002416>
- Hemilä H, Chalker E (2013) Vitamin C for preventing and treating the common cold. *Cochrane Database Syst Rev*. <https://doi.org/10.1002/14651858.CD000980.PUB4>
- Hill C, Guarner F, Reid G et al (2014) Expert consensus document. the international scientific association for probiotics and prebiotics consensus statement on the scope and appropriate use of the term probiotic. *Nat Rev Gastroenterol Hepatol* 11:506–514. <https://doi.org/10.1038/NRGASTRO.2014.66>
- Holland A, Hill CJ, Jones AY, McDonald CF (2012) Breathing exercises for chronic obstructive pulmonary disease. *Cochrane Database Syst Rev* 10:CD008250. <https://doi.org/10.1002/14651858.CD008250.pub2>
- Hopkins SR, Dominelli PB, Davis CK et al (2021) Face masks and the cardiorespiratory response to physical activity in health and disease. *Ann Am Thorac Soc* 18:399–407. <https://doi.org/10.1513/AnnalsATS.202008-990CME>
- Hopkinson NS, Dayer MJ, Moxham J, Polkey MI (2010) Abdominal muscle fatigue following exercise in chronic obstructive pulmonary disease. *Respir Res* 11:1–7. <https://doi.org/10.1186/1465-9921-11-15>
- Hosseini B, Berthon BS, Wark P, Wood LG (2017) Effects of fruit and vegetable consumption on risk of asthma, wheezing and immune responses: a systematic review and meta-analysis. *Nutrients* 9:1–26. <https://doi.org/10.3390/nu9040341>
- Hostetter K, McClaran SR, Cox DG, Dallam G (2016) Triathlete adapts to breathing restricted to the nasal passage without loss in VO₂max or vVO₂max. *J Sport Human Perform* 4:1–7
- Illi SK, Held U, Frank I, Spengler CM (2012) Effect of respiratory muscle training on exercise performance in healthy individuals. *Sports Med* 42:707–724. <https://doi.org/10.1007/BF03262290>
- Jansang S, Mickleborough T, Suksom D (2016) Effects of pursed-lip breathing exercise using windmill toy on lung function and respiratory muscle strength in the elderly. *J Med Assoc Thai* 99:1046–1051
- Janssens W, Mathieu C, Boonen S, Decramer M (2011) Vitamin D deficiency and chronic obstructive pulmonary disease: a vicious circle. *Vitam Horm* 86:379–399. <https://doi.org/10.1016/B978-0-12-386960-9.00017-4>
- Jat KR, Goel N, Gupta N et al (2021) Efficacy of vitamin D supplementation in asthmatic children with vitamin D deficiency: a randomized controlled trial (ESDAC trial). *Pediatr Allergy Immunol* 32:479–488. <https://doi.org/10.1111/PAI.13415>
- Jensen D, Ofir D, O'Donnell DE (2009) Effects of pregnancy, obesity and aging on the intensity of perceived breathlessness during exercise in healthy humans. *Respir Physiol Neurobiol* 167:87–100. <https://doi.org/10.1016/J.RESP.2009.01.011>
- Johnson BD, Babcock MA, Suman OE, Dempsey JA (1993) Exercise-induced diaphragm fatigue in healthy humans. *J Physiol* 460:385–405. <https://doi.org/10.1113/jphysiol.1993.sp019477>
- Jolliffe DA, Greenberg L, Hooper RL et al (2017) Vitamin D supplementation to prevent asthma exacerbations: a systematic review and meta-analysis of individual participant data. *Lancet Respir Med* 5:881–890. [https://doi.org/10.1016/S2213-2600\(17\)30306-5](https://doi.org/10.1016/S2213-2600(17)30306-5)
- Jolliffe DA, Greenberg L, Hooper RL et al (2019) Vitamin D to prevent exacerbations of COPD: systematic review and meta-analysis of individual participant data from randomised controlled trials. *Thorax* 74:337–345. <https://doi.org/10.1136/THORA.XJNL-2018-212092>
- Jolliffe DA, Camargo CA, Sluyter JD et al (2021) Vitamin D supplementation to prevent acute respiratory infections: a systematic review and meta-analysis of aggregate data from randomised controlled trials. *Lancet Diabetes Endocrinol* 9:276–292. [https://doi.org/10.1016/S2213-8587\(21\)00051-6](https://doi.org/10.1016/S2213-8587(21)00051-6)
- Kacmarek RM, Ripple R, Cockrill BA et al (1996) Inhaled nitric oxide. A bronchodilator in mild asthmatics with methacholine-induced bronchospasm. *Am J Respir Crit Care Med* 153:128–135. <https://doi.org/10.1164/AJRCCM.153.1.8542105>
- Kanezaki M, Ebihara S (2017) Effect of the cooling sensation induced by olfactory stimulation by L-menthol on dyspnoea: a pilot study. *Eur Respir J* 49:1601823. <https://doi.org/10.1183/13993003.01823-2016>
- Kanezaki M, Terada K, Ebihara S (2020) Effect of olfactory stimulation by L-menthol on laboratory-induced dyspnea in COPD. *Chest* 157:1455–1465. <https://doi.org/10.1016/j.chest.2019.12.028>
- Kanezaki M, Terada K, Ebihara S (2021) L-menthol - a new treatment for breathlessness? *Curr Opin Support Palliat Care* 15:233–238. <https://doi.org/10.1097/SPC.0000000000000569>

- Kau AL, Ahern PP, Griffin NW et al (2011) Human nutrition, the gut microbiome and the immune system. *Nature* 474:327–336. <https://doi.org/10.1038/NATURE10213>
- Knudson D (2009) Significant and meaningful effects in sports biomechanics research. *Sports Biomech* 8:96–104. <https://doi.org/10.1080/14763140802629966>
- Köteles F, Babulka P, Szemerszky R et al (2018) Inhaled peppermint, rosemary and eucalyptus essential oils do not change spirometry in healthy individuals. *Physiol Behav* 194:319–323. <https://doi.org/10.1016/j.physbeh.2018.06.022>
- Krakow B, Melendrez D, Sisley B et al (2006) Nasal dilator strip therapy for chronic sleep-maintenance insomnia and symptoms of sleep-disordered breathing: a randomized controlled trial. *Sleep* 29:16–28. <https://doi.org/10.1007/s11325-005-0037-7>
- Kumar A, Mastana SS, Lindley MR (2016) EPA/DHA dietary supplementation attenuates exercise-induced bronchoconstriction in physically active asthmatic males. *Cogent Med*. <https://doi.org/10.1080/2331205X.2016.1172696>
- Kumar J, Kumar P, Goyal JP et al (2021) Vitamin D supplementation in childhood asthma: a systematic review and meta-analysis of randomised controlled trials. *ERJ Open Res* 8:00662–02021. <https://doi.org/10.1183/23120541.00662-2021>
- Laborde S, Allen MS, Borges U et al (2022) Effects of voluntary slow breathing on heart rate and heart rate variability: a systematic review and a meta-analysis. *Neurosci Biobehav Rev* 138:104711. <https://doi.org/10.1016/j.neubiorev.2022.104711>
- LaComb CO, Tandy RD, Ping Lee S et al (2017) Oral versus nasal breathing during moderate to high intensity submaximal aerobic exercise. *Int J Kinesiol Sports Sci* 5:8. <https://doi.org/10.7575/aiac.ijkss.v.5n.1p.8>
- Langer D, Jacome C, Charusisin N et al (2013) Measurement validity of an electronic inspiratory loading device during a loaded breathing task in patients with COPD. *Respir Med* 107:633–635. <https://doi.org/10.1016/j.rmed.2013.01.020>
- Lareau SC, Fahy B, Meek P (2020) Breathlessness – shortness of breath. *Am J Respir Crit Care Med* 202:3–4
- Larsen KS, Skoffer B, Gregersen Oestergaard L et al (2020) The effects of various respiratory physiotherapies after lung resection: a systematic review. *Physiother Theory Pract* 36:1201–1219. <https://doi.org/10.1080/09593985.2018.1564095>
- Lehoucq A, Mathieu C, Carremans C et al (2012) High doses of vitamin D to reduce exacerbations in chronic obstructive pulmonary disease: a randomized trial. *Ann Intern Med* 156:105–120. <https://doi.org/10.7326/0003-4819-156-2-201201170-00004>
- Leith DE, Bradley M (1976) Ventilatory muscle strength and endurance training. *J Appl Physiol* 41:508–516. <https://doi.org/10.1152/JAPPL.1976.41.4.508>
- Lemoine C, Brigham E, Woo H et al (2020) Relationship between omega-3 and omega-6 fatty acid intake and chronic obstructive pulmonary disease morbidity. *Ann Am Thorac Soc* 17:378–381. <https://doi.org/10.1513/ANNALSATS.201910-740RL>
- Liu J, Dong YQ, Yin J et al (2019) Meta-analysis of vitamin D and lung function in patients with asthma. *Respir Res* 20:1–11. <https://doi.org/10.1186/S12931-019-1072-4>
- Lorca-Santiago J, Jiménez SL, Pareja-Galeano H, Lorenzo A (2020) Inspiratory muscle training in intermittent sports modalities: a systematic review. *Int J Environ Res Public Health* 17:4448. <https://doi.org/10.3390/IJERPH17124448>
- Lu Y, Li P, Li N et al (2020) Effects of home-based breathing exercises in subjects with COPD. *Respir Care* 65:377–387. <https://doi.org/10.4187/respcare.07121>
- Lundberg JON, Settergren G, Gelinder S et al (1996) Inhalation of nasally derived nitric oxide modulates pulmonary function in humans. *Acta Physiol Scand* 158:343–347. <https://doi.org/10.1046/j.1365-201X.1996.557321000.x>
- Luo C, Sun Y, Zeng Z et al (2022) Vitamin D supplementation in pregnant women or infants for preventing allergic diseases: a systematic review and meta-analysis of randomized controlled trials. *Chin Med J* 135:276–284. <https://doi.org/10.1097/CM9.0000000000001951>
- Macêdo TMF, Freitas DA, Chaves GSS et al (2016) Breathing exercises for children with asthma. *Cochrane Database Syst Rev*. <https://doi.org/10.1002/14651858.CD011017>
- Mai X-M, Langhammer A, Camargo CA, Chen Y (2012) Original contribution serum 25-hydroxyvitamin D levels and incident asthma in adults: the HUNT Study. *Am J Epidemiol* 176:1169–1176. <https://doi.org/10.1093/aje/kws235>
- Mangla PK, Menon MPS (1981) Effect of nasal and oral breathing on exercise-induced asthma. *Clin Exp Allergy* 11:433–439. <https://doi.org/10.1111/J.1365-2222.1981.TB01616.X>
- Manifield J, Winnard A, Hume E et al (2021) Inspiratory muscle training for improving inspiratory muscle strength and functional capacity in older adults: a systematic review and meta-analysis. *Age Ageing* 50:716–724. <https://doi.org/10.1093/AGEING/AFAA221>
- Manousaki D, Paternoster L, Standl M et al (2017) Vitamin D levels and susceptibility to asthma, elevated immunoglobulin E levels, and atopic dermatitis: a Mendelian randomization study. *PLoS Med* 14:e1002294. <https://doi.org/10.1371/journal.pmed.1002294>
- Marciniuk DD, Goodridge D, Hernandez P et al (2011) Managing dyspnea in patients with advanced chronic obstructive pulmonary disease: a Canadian thoracic society clinical practice guideline. *Can Respir J* 18:69–78. <https://doi.org/10.1155/2011/745047>
- Marotta N, Demeco A, Moggio L et al (2020) Comparative effectiveness of breathing exercises in patients with chronic obstructive pulmonary disease. *Complementary Ther Clin Pract* 41:101260. <https://doi.org/10.1016/j.ctcp.2020.101260>
- Marsland BJ, Trompette A, Gollwitzer ES (2015) The gut-lung axis in respiratory disease. *Ann Am Thorac Soc* 12:150–156. <https://doi.org/10.1513/ANNALSATS.201503-133AW>
- Martineau AR, Jolliffe DA, Hooper RL et al (2017) Vitamin D supplementation to prevent acute respiratory tract infections: systematic review and meta-analysis of individual participant data. *Br Med J* 356:i6583. <https://doi.org/10.1136/BMJ.I6583>
- Mason SA, Trewin AJ, Parker L, Wadley GD (2020) Antioxidant supplements and endurance exercise: current evidence and mechanistic insights. *Redox Biol* 35:101471. <https://doi.org/10.1016/J.REDOX.2020.101471>
- McConnell AK (2012) CrossTalk opposing view: respiratory muscle training does improve exercise tolerance. *J Physiol* 590:3397–3398. <https://doi.org/10.1113/JPHYSIOL.2012.235572>
- McConnell AK, Lomax M (2006) The influence of inspiratory muscle work history and specific inspiratory muscle training upon human limb muscle fatigue. *J Physiol* 577:445–457. <https://doi.org/10.1113/jphysiol.2006.117614>
- McConnell AK, Romer LM (2004a) Dyspnoea in health and obstructive pulmonary disease: the role of respiratory muscle function and training. *Sports Med* 34:117–132. <https://doi.org/10.2165/00007256-200434020-00005>
- McConnell AK, Romer LM (2004b) Respiratory muscle training in healthy humans: resolving the controversy. *Int J Sports Med* 25:284–293. <https://doi.org/10.1055/S-2004-815827/ID/38>
- McNarry MA, Berg RMG, Shelley J et al (2022) Inspiratory muscle training enhances recovery post COVID-19: a randomised controlled trial. *Eur Respir J*. <https://doi.org/10.1183/13993003.03101-2021>
- Mendes LP, Moraes KS, Hoffman M et al (2019) Effects of diaphragmatic breathing with and without pursed-lips breathing in

- subjects with COPD. *Respir Care* 64:136–144. <https://doi.org/10.4187/respcare.06319>
- Mickleborough TD, Lindley MR (2014) The effect of combining fish oil and vitamin C on airway inflammation and hyperpnea-induced bronchoconstriction in asthma. *J Allergy Ther* 5:1–10. <https://doi.org/10.4172/2155-6121.1000184>
- Mickleborough TD, Murray RL, Ionescu AA, Lindley MR (2003) Fish oil supplementation reduces severity of exercise-induced bronchoconstriction in elite athletes. *Am J Respir Crit Care Med* 168:1181–1189. <https://doi.org/10.1164/RCCM.200303-373OC>
- Mickleborough TD, Lindley MR, Ionescu AA, Fly AD (2006) Protective effect of fish oil supplementation on exercise-induced bronchoconstriction in asthma. *Chest* 129:39–49. <https://doi.org/10.1378/CHEST.129.1.39>
- Millqvist E, Ternesten-Hasséus E, Bende M (2013) Inhalation of menthol reduces capsaicin cough sensitivity and influences inspiratory flows in chronic cough. *Respir Med* 107:433–438. <https://doi.org/10.1016/j.rmed.2012.11.017>
- Molgat-Seon Y, Dominelli PB, Ramsook AH et al (2018) The effects of age and sex on mechanical ventilatory constraint and dyspnea during exercise in healthy humans. *J Appl Physiol* 124:1092–1106. <https://doi.org/10.1152/jappphysiol.00608.2017>
- Morris GN, Rich GF (1997) Inhaled nitric oxide as a selective pulmonary vasodilator in clinical anesthesia. *Am Assoc Nurse Anesthesiol* 65:59–67
- Morton AR, King K, Papalia S et al (1995) Comparison of maximal oxygen consumption with oral and nasal breathing. *Aust J Sci Med Sport* 27:51–55
- Mourgeon E, Puybasset L, Law-Koune JD et al (1997) Inhaled nitric oxide in acute respiratory distress syndrome with and without septic shock requiring norepinephrine administration: a dose-response study. *Crit Care* 1:25–39. <https://doi.org/10.1186/cc4>
- Mündel T, Jones DA (2010) The effects of swilling an L(-)-menthol solution during exercise in the heat. *Eur J Appl Physiol* 109:59–65. <https://doi.org/10.1007/S00421-009-1180-9>
- Naclerio RM, Pinto J, Assanasen P, Baroody FM (2007) Observations on the ability of the nose to warm and humidify inspired air. *Rhinology* 45:102–111
- Naclerio RM, Pinto J, Assanasen P, Baroody FM (2007) Observations on the ability of the nose to warm and humidify inspired air. *Rhinology* 45:102–111
- Nguyen JD, Duong H (2021) Pursed-lip breathing. StatPearls Publishing, Treasure Island, Florida, US
- Nield MA, Soo Hoo GW, Roper JM, Santiago S (2007) Efficacy of pursed-lips breathing: a breathing pattern retraining strategy for dyspnea reduction. *J Cardiopulm Rehabil Prev* 27:237–244. <https://doi.org/10.1097/01.HCR.0000281770.82652.CB>
- Niinimaa V, Cole P, Mintz S, Shepard RJ (1980) The switching point from nasal to oronasal breathing. *Respir Physiol* 42:61–71. [https://doi.org/10.1016/0034-5687\(80\)90104-8](https://doi.org/10.1016/0034-5687(80)90104-8)
- Nishino T, Tagaito Y, Sakurai Y (1997) Nasal inhalation of l-menthol reduces respiratory discomfort associated with loaded breathing. *Am J Respir Crit Care Med* 156:309–313. <https://doi.org/10.1164/ajrccm.156.1.9609059>
- Nitzan I, Mimouni FB, Bin NA et al (2022) Vitamin D and asthma: a systematic review of clinical trials. *Curr Nutr Rep* 11:311–317. <https://doi.org/10.1007/S13668-022-00411-6>
- Noss MJ, Ciesla R, Shanga G (2019) Sleep quality and congestion with breathe right nasal strips: two randomized controlled trials. *Adv Ther* 36:1975–1985. <https://doi.org/10.1007/s12325-019-01005-5>
- O’Kroy JA, James T, Miller JM et al (2001) Effects of an external nasal dilator on the work of breathing during exercise. *Med Sci Sports Exerc* 33:454–458. <https://doi.org/10.1097/00005768-200103000-00019>
- O’Neill B, Mac MJ, Bradley J (2006) Short-burst oxygen therapy in chronic obstructive pulmonary disease. *Respir Med* 100:1129–1138. <https://doi.org/10.1016/J.RMED.2006.03.038>
- Parisien-La Salle S, Abel Rivest E, Boucher VG et al (2019) Effects of pursed lip breathing on exercise capacity and dyspnea in patients with interstitial lung disease: a randomized, crossover study. *J Cardiopulm Rehabil Prev* 39:112–117. <https://doi.org/10.1097/HCR.0000000000000387>
- Parvzian MK, Dhaliwal M, Li J et al (2020) Relationship between dietary patterns and COPD: a systematic review and meta-analysis. *ERJ Open Res* 6:00168–02019. <https://doi.org/10.1183/23120541.00168-2019>
- Pasquina P, Tramèr MR, Granier JM, Walder B (2006) Respiratory physiotherapy to prevent pulmonary complications after abdominal surgery: a systematic review. *Chest* 130:1887–1899. <https://doi.org/10.1378/CHEST.130.6.1887>
- Patel MS, Hart N, Polkey MI (2012) CrossTalk proposal: training the respiratory muscles does not improve exercise tolerance. *J Physiol* 590:3393–3395. <https://doi.org/10.1113/JPHYSIOL.2012.235408>
- Pereira EJ, Sim L, Driver HS et al (2013) The effect of inhaled menthol on upper airway resistance in humans: a randomized controlled crossover study. *Can Respir J* 20:e1-4. <https://doi.org/10.1155/2013/383019>
- Peters CM, Leahy MG, Hohert G et al (2021) Airway luminal area and the resistive work of breathing during exercise in healthy young females and males. *J Appl Physiol* 131:1750–1761. <https://doi.org/10.1152/JAPPLPHYSIOL.00418.2021>
- Pham H, Rahman A, Majidi A et al (2019) Acute respiratory tract infection and 25-hydroxyvitamin D concentration: a systematic review and meta-analysis. *Environ Res Public Health* 16:3020. <https://doi.org/10.3390/ijerph16173020>
- Philips DB, Neder JA, Elbehairy AF et al (2021) Qualitative components of dyspnea during incremental exercise across the COPD continuum. *Med Sci Sports Exerc* 53:2467–2476. <https://doi.org/10.1249/MSS.00000000000002741>
- Porcari JP, Probst L, Forrester K et al (2016) Effect of wearing the elevation training mask on aerobic capacity, lung function, and hematological variables. *J Sports Sci Med* 15:375–386
- Rashidi K, Razi B, Darand M et al (2021) Effect of probiotic fermented dairy products on incidence of respiratory tract infections: a systematic review and meta-analysis of randomized clinical trials. *Nutr J* 20:61. <https://doi.org/10.1186/S12937-021-00718-0>
- Ricciardolo FLM (2003) Multiple roles of nitric oxide in the airways. *Thorax* 58:175–182. <https://doi.org/10.1136/thorax.58.2.175>
- Ries AL, Bauldoff GS, Carlin BW et al (2007) Pulmonary rehabilitation: joint ACCP/AACVPR evidence-based clinical practice guidelines. *Chest* 131:S4–S42. <https://doi.org/10.1378/CHEST.06-2418>
- Roberfroid M, Gibson GR, Hoyle L et al (2010) Prebiotic effects: metabolic and health benefits. *Br J Nutr* 104:S1–S6. <https://doi.org/10.1017/S0007114510003363>
- Rodrigues GD, Gurgel JL, da Silva Soares PP (2020) Inspiratory muscles contributions to postural control in healthy older women. *Respir Physiol Neurobiol* 281:103505. <https://doi.org/10.1016/J.RESP.2020.103505>
- Romer LM, McConnell AK (2003) Specificity and reversibility of inspiratory muscle training. *Med Sci Sports Exerc* 35:237–244. <https://doi.org/10.1249/01.MSS.0000048642.58419.1E>
- Ruuskanen O, Luoto R, Valtonen M et al (2022) Respiratory viral infections in athletes: many unanswered questions. *Sports Med* 52:2013–2021. <https://doi.org/10.1007/S40279-022-01660-9>
- Sadan O, Shushan S, Eldar I et al (2005) The effects of an external nasal dilator on labor. *Am J Rhinol* 19:221–224
- Sagar S, Morgan ME, Chen S et al (2014) Bifidobacterium breve and *Lactobacillus rhamnosus* treatment is as effective as budesonide

- at reducing inflammation in a murine model for chronic asthma. *Respir Res* 15:46. <https://doi.org/10.1186/1465-9921-15-46>
- Sandnes A, Andersen T, Clemm HH et al (2022) Clinical responses following inspiratory muscle training in exercise-induced laryngeal obstruction. *Eur Arch Otorhinolaryngol* 279:2511–2522. <https://doi.org/10.1007/S00405-021-07214-5>
- Santino TA, Chaves GSS, Freitas DA et al (2020) Breathing exercises for adults with asthma. *Cochrane Database Syst Rev*. <https://doi.org/10.1002/14651858.CD001277.pub4>
- Schaeffer MR, McBride E, Mitchell RA et al (2021) Effects of the Turbine™ on ventilatory and sensory responses to incremental cycling. *Med Sci Sports Exerc* 53:192–199. <https://doi.org/10.1249/MSS.0000000000002427>
- Schultz K, Jelusic D, Wittmann M et al (2018) Inspiratory muscle training does not improve clinical outcomes in 3-week COPD rehabilitation: results from a randomised controlled trial. *Eur Respir J* 51:1702000. <https://doi.org/10.1183/13993003.02000-2017>
- Seixas MB, Almeida LB, Trevizan PF et al (2020) Effects of inspiratory muscle training in older adults. *Respir Care* 65:535–544. <https://doi.org/10.4187/RESPCARE.06945>
- Settergren G, Angdin M, Astudillo R et al (1998) Decreased pulmonary vascular resistance during nasal breathing: modulation by endogenous nitric oxide from the paranasal sinuses. *Acta Physiol Scand* 163:235–239. <https://doi.org/10.1046/j.1365-201x.1998.00352.x>
- Sheel AW, Boushel R, Dempsey JA (2018) Competition for blood flow distribution between respiratory and locomotor muscles: implications for muscle fatigue. *J Appl Physiol* 125:820–831. <https://doi.org/10.1152/jappphysiol.00189.2018>
- Shturman-Ellstein R, Zeballos RJ, Buckley JM, Souhrada JF (1978) The beneficial effect of nasal breathing on exercise-induced bronchoconstriction. *Am Rev Respir Dis* 118:65–73. <https://doi.org/10.1164/ARRD.1978.118.1.65>
- Sluyter JD, Camargo CA, Waayer D et al (2017) Effect of monthly, high-dose, long-term vitamin D on lung function: a randomized controlled trial. *Nutrients* 9:1353. <https://doi.org/10.3390/NU9121353>
- Spahija J, De Marchie M, Grassino A (2005) Effects of imposed pursed-lips breathing on respiratory mechanics and dyspnea at rest and during exercise in COPD. *Chest* 128:640–650. <https://doi.org/10.1378/chest.128.2.640>
- Spruit MA, Singh SJ, Garvey C et al (2013) An official American thoracic society/European respiratory society statement: key concepts and advances in pulmonary rehabilitation. *Am J Respir Crit Care Med* 188:e13–e64. <https://doi.org/10.1164/RCCM.201309-1634ST>
- Stanford G, Ryan H, Solis-Moya A (2020) Respiratory muscle training for cystic fibrosis. *Cochrane Database Syst Rev* 12:CD006112. <https://doi.org/10.1002/14651858.CD006112.PUB5>
- Stoodley I, Garg M, Scott H et al (2019) Higher omega-3 index is associated with better asthma control and lower medication dose: a cross-sectional study. *Nutrients* 12:74. <https://doi.org/10.3390/NU12010074>
- Strasser B, Geiger D, Schauer M et al (2016) Probiotic supplements beneficially affect tryptophan-kynurenine metabolism and reduce the incidence of upper respiratory tract infections in trained athletes: a randomized, double-blinded, placebo-controlled trial. *Nutrients* 8:752. <https://doi.org/10.3390/NU8110752>
- Tamaoki J, Chiyotani A, Sakai A et al (1995) Effect of menthol vapour on airway hyperresponsiveness in patients with mild asthma. *Respir Med* 89:503–504. [https://doi.org/10.1016/0954-6111\(95\)90127-2](https://doi.org/10.1016/0954-6111(95)90127-2)
- Tanaka Y, Morikawa T, Honda Y (1988) An assessment of nasal functions in control of breathing. *J Appl Physiol* 65:1520–1524. <https://doi.org/10.1152/jappl.1988.65.4.1520>
- Taylor BJ, How SC, Romer LM (2006) Exercise-induced abdominal muscle fatigue in healthy humans. *J Appl Physiol* 100:1554–1562. <https://doi.org/10.1152/JAPPLPHYSIOL.01389.2005>
- Tecklenburg SL, Mickleborough TD, Fly AD et al (2007) Ascorbic acid supplementation attenuates exercise-induced bronchoconstriction in patients with asthma. *Respir Med* 101:1770–1778. <https://doi.org/10.1016/J.RMED.2007.02.014>
- Tecklenburg-Lund S, Mickleborough TD, Turner LA et al (2010) Randomized controlled trial of fish oil and montelukast and their combination on airway inflammation and hyperpnea-induced bronchoconstriction. *PLoS One* 5:e13487. <https://doi.org/10.1371/JOURNAL.PONE.0013487>
- Thien FC, De LS, Woods RK, Abramson MJ (2002) Dietary marine fatty acids (fish oil) for asthma in adults and children. *Cochrane Database Syst Rev*. <https://doi.org/10.1002/14651858.CD001283>
- Thoman RL, Stoker GL, Ross JC (1966) The efficacy of pursed-lips breathing in patients with chronic obstructive pulmonary disease. *Am Rev Respir Dis* 93:100–106. <https://doi.org/10.1164/arrd.1966.93.1.100>
- Thomas DQ, Larson BM, Rahija MR, McCaw ST (2001) Nasal strips do not affect cardiorespiratory measures during recovery from anaerobic exercise. *J Strength Cond Res* 15:341–343
- Tiep BL, Burns M, Kao D et al (1986) Pursed lips breathing training using ear oximetry. *Chest* 90:218–221. <https://doi.org/10.1378/chest.90.2.218>
- Tiller NB, Campbell IG, Romer LM (2017) Influence of upper-body exercise on the fatigability of human respiratory muscles. *Med Sci Sports Exerc* 49:1461–1472. <https://doi.org/10.1249/MSS.0000000000001251>
- Tiller NB, Sullivan JP, Ekkekakis P (2022) Baseless claims and pseudoscience in health and wellness: a call to action for the sports, exercise, and nutrition-science community. *Sports Med* 2022:1–5. <https://doi.org/10.1007/S40279-022-01702-2>
- Törnberg DCF, Marteus H, Schedin U et al (2002) Nasal and oral contribution to inhaled and exhaled nitric oxide: a study in tracheotomized patients. *Eur Respir J* 19:859–864. <https://doi.org/10.1183/09031936.02.00273502>
- Torres-Castro R, Solis-Navarro L, Puppo H et al (2022) Respiratory muscle training in patients with obstructive sleep apnoea: a systematic review and meta-analysis. *Clocks Sleep* 4:219–229. <https://doi.org/10.3390/CLOCKSSLEEP4020020>
- Tounsi B, Acheche A, Lelard T et al (2021) Effects of specific inspiratory muscle training combined with whole-body endurance training program on balance in COPD patients: randomized controlled trial. *PLoS One* 16:e0257595. <https://doi.org/10.1371/JOURNAL.PONE.0257595>
- Tran D, Munoz P, Lau EMT et al (2021) Inspiratory muscle training improves inspiratory muscle strength and functional exercise capacity in pulmonary arterial hypertension and chronic thromboembolic pulmonary hypertension: a pilot randomised controlled study. *Heart Lung Circ* 30:388–395. <https://doi.org/10.1016/j.hlc.2020.06.006>
- Ubolnuar N, Tantisawat A, Thaveeratitham P et al (2019) Effects of breathing exercises in patients with chronic obstructive pulmonary disease: systematic review and meta-analysis. *Ann Rehabil Med* 43:509–523. <https://doi.org/10.5535/arm.2019.43.4.509>
- van der Schans CP, de Jong W, de Vries G et al (1997) Respiratory muscle activity and pulmonary function during acutely induced airways obstruction. *Physiother Res Int* 2:167–177. <https://doi.org/10.1002/pri.98>
- Varraso R, Barr RG, Willett WC et al (2015) Fish intake and risk of chronic obstructive pulmonary disease in 2 large US cohorts.

- Am J Clin Nutr 101:354–361. <https://doi.org/10.3945/AJCN.114.094516>
- Venter C, Eyerich S, Sarin T, Klatt KC (2020) Nutrition and the immune system: a complicated tango. *Nutrients* 12:818. <https://doi.org/10.3390/nu12030818>
- Verges S, Lenherr O, Haner AC et al (2007) Increased fatigue resistance of respiratory muscles during exercise after respiratory muscle endurance training. *Am J Physiol Regul Integr Comp Physiol* 292:1246–1253. <https://doi.org/10.1152/ajpregu.00409.2006>
- Verges S, Renggli AS, Notter DA, Spengler CM (2009) Effects of different respiratory muscle training regimes on fatigue-related variables during volitional hyperpnoea. *Respir Physiol Neurobiol* 169:282–290. <https://doi.org/10.1016/J.RESP.2009.09.005>
- Verheijden KAT, Willemsen LEM, Braber S et al (2015) Dietary galacto-oligosaccharides prevent airway eosinophilia and hyperresponsiveness in a murine house dust mite-induced asthma model. *Respir Res* 16:17. <https://doi.org/10.1186/S12931-015-0171-0>
- Verheijden KAT, Willemsen LEM, Braber S et al (2016) The development of allergic inflammation in a murine house dust mite asthma model is suppressed by synbiotic mixtures of non-digestible oligosaccharides and *Bifidobacterium breve* M-16V. *Eur J Nutr* 55:1141–1151. <https://doi.org/10.1007/S00394-015-0928-8>
- Vermoen CJ, Verbraak AF, Bogaard JM (1998) Effect of a nasal dilator on nasal patency during normal and forced nasal breathing. *Int J Sports Med* 19:109–113. <https://doi.org/10.1055/s-2007-971891>
- Vlieg-Boerstra B, De Jong N, I, Meyer R, et al (2021) Nutrient supplementation for prevention of viral respiratory tract infections in health subjects: a systematic review and meta-analysis. *Allergy* 77:1373–1388. <https://doi.org/10.1111/all.15136>
- WADA (2022) Prohibited methods (prohibited at all times). *World Anti-Doping Code Prohib List* 2022:13
- Walsh NP (2019) Nutrition and athlete immune health: new perspectives on an old paradigm. *Sports Med* 49:S153–S168. <https://doi.org/10.1007/S40279-019-01160-3>
- Wenzel M, Schönhofer B, Siemon K, Köhler D (1997) Nasal strips without effect on obstructive sleep apnea and snoring. *Pneumologie* 51:1108–1110
- West NP, Pyne DB, Cripps AW et al (2011) *Lactobacillus fermentum* (PCC®) supplementation and gastrointestinal and respiratory-tract illness symptoms: a randomised control trial in athletes. *Nutr J* 10:30. <https://doi.org/10.1186/1475-2891-10-30>
- West NP, Horn PL, Pyne DB et al (2014) Probiotic supplementation for respiratory and gastrointestinal illness symptoms in healthy physically active individuals. *Clin Nutr* 33:581–587. <https://doi.org/10.1016/J.CLNU.2013.10.002>
- Williams NC, Johnson MA, Shaw DE et al (2016) A prebiotic galacto-oligosaccharide mixture reduces severity of hyperpnoea-induced bronchoconstriction and markers of airway inflammation. *Br J Nutr* 116:798–804. <https://doi.org/10.1017/S0007114516002762>
- Williams NC, Hunter KA, Shaw DE et al (2017) Comparable reductions in hyperpnoea-induced bronchoconstriction and markers of airway inflammation after supplementation with 6·2 and 3·1 g/d of long-chain n-3 PUFA in adults with asthma. *Br J Nutr* 117:1379–1389. <https://doi.org/10.1017/S0007114517001246>
- Witt JD, Guenette JA, Rupert JL et al (2007) Inspiratory muscle training attenuates the human respiratory muscle metaboreflex. *J Physiol* 584:1019–1028. <https://doi.org/10.1113/jphysiol.2007.140855>
- Woods A, Gustafson O, Williams M, Stiger R (2022) The effects of inspiratory muscle training on inspiratory muscle strength, lung function and quality of life in adults with spinal cord injuries: a systematic review and meta-analysis. *Disabil Rehabil*. <https://doi.org/10.1080/09638288.2022.2107085>
- World Health Organization (2022) Chronic obstructive pulmonary disease (COPD) [fact sheet]. [https://www.who.int/news-room/fact-sheets/detail/chronic-obstructive-pulmonary-disease-\(copd\)](https://www.who.int/news-room/fact-sheets/detail/chronic-obstructive-pulmonary-disease-(copd)). Accessed 26 Sep 2022
- Worraphan S, Thammata A, Chittawatanarat K et al (2020) Effects of inspiratory muscle training and early mobilization on weaning of mechanical ventilation: a systematic review and network meta-analysis. *Arch Phys Med Rehabil* 101:2002–2014. <https://doi.org/10.1016/J.APMR.2020.07.004>
- Yagihara F, Lorenzi-Filho G, Santos-Silva R (2017) Nasal dilator strip is an effective placebo intervention for severe obstructive sleep apnea. *J Clin Sleep Med* 13:215–221. <https://doi.org/10.5664/jcsm.6450>
- Yang H, Xun P, He K (2013) Fish and fish oil intake in relation to risk of asthma: a systematic review and meta-analysis. *PLoS One* 8:e80048. <https://doi.org/10.1371/JOURNAL.PONE.0080048>
- Yang Y, Wei L, Wang S et al (2022) The effects of pursed lip breathing combined with diaphragmatic breathing on pulmonary function and exercise capacity in patients with COPD: a systematic review and meta-analysis. *Physiother Theory Pract* 38:847–857. <https://doi.org/10.1080/09593985.2020.1805834>
- Yepes-Núñez JJ, Brozek JL, Fiocchi A et al (2018) Vitamin D supplementation in primary allergy prevention: systematic review of randomized and non-randomized studies. *Allergy* 73:37–49. <https://doi.org/10.1111/all.13241>
- Yu H, Su X, Lei T et al (2021) Effect of omega-3 fatty acids on chronic obstructive pulmonary disease: a systematic review and meta-analysis of randomized controlled trials. *Int J Chron Obstruct Pulmon Dis* 16:2677–2686. <https://doi.org/10.2147/COPD.S331154>
- Zaccaro A, Piarulli A, Laurino M et al (2018) How breath-control can change your life: a systematic review on psycho-physiological correlates of slow breathing. *Front Hum Neurosci* 12:1–16. <https://doi.org/10.3389/fnhum.2018.00353>
- Zaki S, Moiz JA, Mujaddadi A et al (2022) Does inspiratory muscle training provide additional benefits during pulmonary rehabilitation in people with interstitial lung disease? A randomized control trial. *Physiother Theory Pract* 13:24–26. <https://doi.org/10.1080/09593985.2021.2024311>
- Zemková E (2014) Significantly and practically meaningful differences in balance research: P values and/or effect sizes? *Sports Med* 44:879–886. <https://doi.org/10.1007/s40279-014-0185-7>
- Zhu M, Wang T, Wang C, Ji Y (2016) The association between vitamin D and COPD risk, severity, and exacerbation: an updated systematic review and meta-analysis. *Int J Chron Obstruct Pulmon Dis* 11:2597–2607. <https://doi.org/10.2147/COPD.S101382>

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Springer Nature or its licensor (e.g. a society or other partner) holds exclusive rights to this article under a publishing agreement with the author(s) or other rightsholder(s); author self-archiving of the accepted manuscript version of this article is solely governed by the terms of such publishing agreement and applicable law.